(19) World Intellectual Property Organization International Bureau



T 1980) BINGRON IN GUNN BRANCH DAN IN HAT ROOM BINGROND BURGE HALL HAN BINGROND BURGE

(43) International Publication Date 1 March 2001 (01.03.2001)

PCT

(10) International Publication Number WO 01/14395 A2

(51) International Patent Classification7:

C07H 15/00

(21) International Application Number: PCT/US00/22894

(22) International Filing Date: 18 August 2000 (18.08.2000)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data: 60/150,088

20 August 1999 (20.08.1999) US

- (71) Applicant: SLOAN-KETTERING INSTITUTE FOR CANCER RESEARCH [US/US]; 1275 York Avenue, New York, NY 10021 (US).
- (72) Inventors: DANISHEFSKY, Samuel, J.; 22 Brayton Street, Englewood, NJ 07631 (US). ALLEN, Jennifer, R.; 5886 North Delaware Street, Indianapolis, IN 46220 (US). RAGUPATHI, Govindaswami; 504 East 81st Street, New York, NY 10028 (US). LIVINGSTON, Philip, O.; 156 East 79th Street, New York, NY 10021 (US).

- (74) Agent: SHAIR, Karoline, K., M.; Choate, Hall & Stewart, Exchange Place, 53 State Street, Boston, MA 02109 (US).
- (81) Designated States (national): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

 Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NOVEL GLYCOCONJUGATES, GLYCOAMINO, ACIDS, INTERMEDIATES THERETO, AND USES THEREOF

(57) Abstract: The present invention provides novel n-alkenyl glycosides and glycoconjugates, n-alkyl glycoamino acids, and methods for the synthesis thereof. In another aspect, the present invention provides novel clustered glycopeptides and methods for the synthesis thereof. In still another aspect, the present invention provides methods for the treatment of cancer, preferably for the prevention of recurrence of cancer, and methods for inducing antibodies in a subject, comprising administering to a subject in need, an effective amount of any of the inventive glycoconjugates as disclosed herein.



11/14395 42

NOVEL GLYCOCONJUGATES, GLYCOAMINO ACIDS, INTERMEDIATES THERETO, AND USES THEREOF

5

10

15

20

25

30

35

Priority Information

This application claims priority under § 119(e) of the United States Code to provisional application 60/150,088, filed August 20, 1999, entitled "Synthesis and Bioconjugation of the n-Pentenyl Glycoside of the Tumor-Associated Antigen Fucosyl GM1", the entire contents of which are hereby incorporated by reference.

Government Support

The present invention was supported by the National Institutes of Health Grant Numbers: AI16943 and CA28824. Therefore, the government has certain rights in the present invention.

Background of the Invention

The improvement of existing therapeutics and the development of novel therapeutics to treat and/or prolong survival of cancer patients has been the subject of continuing research in the scientific community. Although certain of these efforts have been directed to more "traditional" chemotherapeutics (e.g., Paclitaxel and other small molecule and/or natural product based therapies) that act by killing malignant cancer cells, it has also been a long-standing goal (Lanzavechis, Science, 260, 937-944; Pardoll et al., Curr. Opin. Immunol. 1993, 5, 719-725; Livingston et al., Curr. Opin. Immunol. 1992, 4, 2; Dranoff et al., Proc. Natl. Acad. Sci. USA 1993, 90, 3539; M.H. Taoet et al., Nature, 1993, 362, 755; T. Boon, Int. J. Cancer 1993, 54, 177) to develop an anti-cancer vaccine to induce an anticancer response. Although cancer vaccines have thus far been perceived as a mode of treatment subsequent to the detection of the disease (for example, by providing an enhanced immunological response), it would be most desirable to develop a selective synthetic vaccine that would be able to provide enhanced protection against tumor recurrence and metastasis when the tumor burden has been rendered minimal through surgery, radiation or other chemotherapeutic treatment.

In general, tumor immunotherapy is based on the theory that tumors possess specific antigens that can be recognized when presented to or processed by a properly

trained immune system. The goal for the development of an effective anticancer vaccine is to break the tolerance which the immune system has for these antigens expressed mainly or exclusively by the tumor, by presenting glycoconjugates as versions of immunostimulatory antigens, to induce an effective immune response. In an effort to achieve this goal, identified cancer carbohydrate antigens such as TF, Tn, sTN, KH-1, Ley and Globo-H have been carefully characterized as being over-expressed at the surface of malignant cells in a variety of cancers (breast, colon, prostate, ovarian, liver, small cell lung and adenocarcinomas). In addition, they have been immunocharacterized by monoclonal antibodies and therefore have relevant serological markers available for immunological studies. Such studies have suggested that patients immunized in an adjuvant setting with carbohydrate-based vaccines produce antibodies reactive with human cancer cells, and that the production of such antibodies prohibits tumor recurrence and correlates with a more favorable diagnosis (see, Livingston et al., J. Cancer Res. 1989, 49, 7045; Ragupathi, G. Cancer Immunol. Immunother. 1996, 43, 152). Additionally, the isolation and careful structural identification of specific carbohydrate antigens overexpressed in cancer cells has provided a framework for an attack using carbohydrate-based tumor immunotherapy (For reviews see (a) Hakomori, S.; Zhang, Y. Chem. Biol. 1997, 4, 97; (b) Toyokuni, T.; Singhal, A. K. Chem. Soc. Rev. 1995, 24, 23 and references therein).

20

25

30

35

A major drawback in using carbohydrate epitopes, however, is that they are generally not readily available by isolation from natural sources. For example, the immense difficulties associated with their purification from natural sources render them virtually nonavailable as homogeneous starting materials for a clinical program. Thus, the incorporation of these naturally occurring epitopes into carrier proteins or any favorable molecular context via conjugation for eliciting a therapeutically useful immunological response is inefficient at best, and often virtually impossible. Therefore, to effectively study these vaccines as therapeutic agents, sufficient material can only be obtained by total chemical synthesis.

In an effort to remedy this problem, one of the continuing research efforts is the development of anti-cancer vaccines that incorporate fully synthetic carbohydrate moieties (For a review, see Danishefsky, S.J.; Allen, J.R. Angew Chem. Int. Ed. 2000, 39,

836-863). One strategy for the development of synthetic anti-cancer vaccines involves the total synthesis of the carbohydrate epitope and its subsequent covalent bioconjugation to carrier protein. The vaccine constructs are then subjected to appropriate mouse immunization studies, with the ultimate goal of advancing to human clinical trials. This strategy has resulted in several fully synthetic tumor associated carbohydrate-based vaccines which are at various stages of advanced pre-clinical and clinical processing. In fact, a Globo-H vaccine is undergoing clinical evaluation for the treatment of prostate and breast carcinomas at the phase II level (see, for example, Ragupathi et al., Angew. Chem. Int. Ed. Engl. 1997, 36, 125) while a Lewis antigen-based vaccine, already tested in ovarian cancer, is awaiting more extensive follow-up evaluation (see, Kudryashov et al. Cancer Immunol. Immunother. 1998, 45, 281).

Although several synthetic constructs have been developed in recent years, as described above, and in other references described herein, there remains a need for the further investigation to develop novel constructs capable of eliciting a more sustained or effective (and preferably selective) immune response. Clearly, in an effort to achieve this goal, it would be useful to develop improved and/or novel synthetic methods to access heretofore synthetically unavailable antigenic components (e.g., more complex antigenic components such as fucosyl GM1, clustered epitopes and similar structures), or to access non-natural structures derived from naturally occurring structures for further immunologic and therapeutic studies.

25

30

35

15

20

Summary of the Invention

In recognition of the need to further develop novel constructs and improved synthetic methods, the present invention, in one aspect, provides novel n-alkenyl glycosides and glycoconjugates, n-alkyl glycoamino acids, and methods for the synthesis thereof. In another aspect, the present invention provides novel clustered glycopeptides and methods for the synthesis thereof. In still another aspect, the present invention provides methods for the treatment of cancer, preferably for the prevention of recurrence of cancer, and methods for inducing antibodies in a subject comprising administering to a subject in need, an effective amount of any of the inventive glycoconjugates as disclosed herein.

10

15

20

The general synthetic methodology disclosed herein involves the realization that the incorporation of an n-alkenyl glycoside protecting group at the reducing end of a carbohydrate acceptor allows for increased coupling efficiencies and accessibility to complex carbohydrates. Thus, the present invention also provides the recognition that for certain protected carbohydrates, the n-alkenyl moieties can serve as useful precursors that can be utilized ultimately for the synthesis of complex glycopeptides.

Thus, in one aspect, the present invention provides novel synthetic methodologies for the synthesis of complex carbohydrates comprising (1) providing a carbohydrate acceptor having a reducing end alkenyl group; (2) providing a suitable donor compound and (3) coupling said donor and acceptor under conditions to generate an alkenyl glycoside. Using this method, complex antigenic alkenyl glycosides are provided, as described above, many of which never before have been provided, which can then be conjugated or further reacted, as described herein, to generate glycoconjugates and glycopeptide structures.

In general, the present invention provides novel compounds and/or conjugates having the general structure:

$$A \sim C(CH_2)_n$$

25

30

wherein R is hydrogen; substituted or unsubstituted alkyl; alkenyl; aryl;
-CH₂CH(CO₂R')(NHR"), wherein R' or R" are each indpendently hydrogen, protecting group, substituted or unsubstituted alkyl, a linker, aryl, peptide, protein or lipid; or NHR", wherein R" is a protein, peptide, or lipid linked to N directly or through a crosslinker; wherein n is 0-8; wherein A is a carbohydrate domain having the structure:

4

15

20

1

$$R_0$$
 R_0 R_0

wherein a, b, c, d, e, f, g, h, i, x, y and z are independently 0, 1, 2 or 3, with the proviso that x, y and z are not simultaneously 0; wherein R₀ is hydrogen, a linear or branched chain alkyl, acyl, arylalkyl or aryl group; wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈ and R₉ are each independently hydrogen, OH, OR¹, NH₂, NHCOR¹, F, CH₂OH, CH₂OR¹, a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein R¹ is hydrogen, CHO, COOR¹¹, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group or a saccharide moiety having the structure:

$$\begin{array}{c|c} R_0 & \begin{array}{c} Z & \begin{array}{c} \\ \end{array} & \end{array} & \begin{array}{c} \\ \end{array} & \begin{array}{c}$$

wherein Y and Z are independently NH or O; wherein k, l, r, s, t, u, v and w are each independently 0, 1 or 2; wherein R₁₀, R₁₁, R₁₂, R₁₃, R₁₄ and R₁₅ are each independently hydrogen, OH, ORⁱⁱⁱ, NH₂, NHCORⁱⁱⁱ, F, CH₂OH, CH₂ORⁱⁱⁱ, or a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein R₁₆ is

(

hydrogen, COOH, COORⁱⁱ, CONHRⁱⁱ, a substituted or unsubstituted linear or branched chain alkyl or aryl group; wherein Rⁱⁱⁱ is hydrogen, CHO, COOR^{iv}, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group; and wherein Rⁱⁱ and R^{iv} are each independently H, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group;

with the proviso that if A is KH-1, N3, globo-H, glycophorin, Tn, TF, STN, (2,3)ST, 2,6-STn or Le^y, and A is α -O-linked, then n is at least 1.

In certain preferred embodiments of the present invention, R is allyl, n is 2 and thus the inventive compound is a n-pentenyl moiety. In certain other embodiments of the present invention, R is NHR'", and the protein R'" is KLH or Bovine Serine Albumin. In still other embodiments of the present invention, R is NHR'", and the lipid R'" is PamCys. It will be appreciated that the protein or lipid can be linked to N directly or through a crosslinker, and thus R'" incorporates proteins, peptides, and lipids, as well as (crosslinker-protein), (crosslinker-peptide) and (crosslinker-lipid) moieties. In certain preferred embodiments, the crosslinker is MMCCH (4-(maleimidomethyl) cyclohexane-1-carboxyl hydrazide).

In other embodiments, the inventive compound is a glycoamino acid and thus R is $CH_2CH(CO_2R')(NHR'')$, which compound has the structure

25

30

10

15

20

In certain preferred embodiments, the glycoamino acids of the present invention are derived from n-pentenyl glycosides and thus n is 3. In certain other preferred embodiments, R' and R" are each a protecting group independently selected from the group consisting of t-butyl, TSE (2-(trimethylsilyl ethyl), Ac (acetyl), Boc (t-butoxycarbonyl), and Fmoc (9-fluoroenyl methoxy carbonyl).

For each of the compounds described above, in certain preferred embodiments the carbohydrate determinant is selected from the group consisting of Globo-H, fucosyl GM1, KH-1, glycophorin, N3, Tn, TF, STN, (2,3)ST, 2,6-STn, and Le^y. In other

15

20

25

30

preferred embodiments, the carbohydrate determinant of the compound contains a Globo-H determinant or fucosyl GM1 determinant, as all or part of the carbohydrate determinant A.

Additionally, the present invention provides methods for the synthesis of novel nalkyl glycoamino acids, as described in more detail below for Globo-H and fucosyl GM1 and their subsequent use to generate novel glycopeptides and synthetic constructs thereof.

In general, the inventive method for the production of these novel glycoamino acids comprises: 1) providing an alkenyl glycosidic moiety, as described herein; 2) subjecting said alkenyl glycosidic moiety to oxidative conditions to generate an aldehyde; 3) subjecting said aldehyde to olefination conditions to generate an enamide ester; 4) subjecting said resulting enamide ester to conditions sufficient to hydrogenate said enamide ester to generate a protected glycoamino acid and 5) deprotecting said protected glycoamino acid under suitable conditions to generate a desired glycoamino acid.

In particular, a method for the synthesis of a glycoamino acid, the structure of which is set forth herein, is provided, which comprises the steps of:

(a) providing an alkenyl glycoside having the structure:

(b) reacting said alkenyl glycoside under suitable conditions to generate an enamideester having the structure:

(b) reacting said enamide ester under suitable conditions to generate a glycoamino acid having the structure:

wherein, for each of the structures above, n is 0-8, wherein A is a carbohydrate domain having the structure:

wherein a, b, c, d, e, f, g, h, i, x, y and z are independently 0, 1, 2 or 3, with the proviso that x, y and z are not simultaneously 0; wherein R₀ is hydrogen, a linear or branched chain alkyl, acyl, arylalkyl or aryl group; wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈ and R₉ are each independently hydrogen, OH, ORⁱ, NH₂, NHCORⁱ, F, CH₂OH, CH₂ORⁱ, a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein Rⁱ is hydrogen, CHO, COORⁱⁱ, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group or a saccharide moiety having the structure:

20

· 5

10

15

wherein Y and Z are independently NH or O; wherein k, l, r, s, t, u, v and w are each independently 0, 1 or 2; wherein R₁₀, R₁₁, R₁₂, R₁₃, R₁₄ and R₁₅ are each independently hydrogen, OH, ORⁱⁱⁱ, NH₂, NHCORⁱⁱⁱ, F, CH₂OH, CH₂ORⁱⁱⁱ, or a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein R₁₆ is hydrogen, COOH, COORⁱⁱ, CONHRⁱⁱ, a substituted or unsubstituted linear or branched chain alkyl or aryl group; wherein Rⁱⁱⁱ is hydrogen, CHO, COOR^{iv}, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group; and wherein Rⁱⁱ and R^{iv} are each independently H, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group;

5

10

15

20

25

30

and wherein for the glycoamino acid structure R' and R" are each independently hydrogen, protecting group, substituted or unsubstituted alkyl, linker, aryl, peptide, protein or lipid; or NHR", wherein R" is a protein, peptide, or lipid, linked to N directly or through a crosslinker. In preferred embodiments, R' and R" are each independently hydrogen or a protecting group. In particularly preferred embodiments R" is a nitrogen protecting group, including, but not limited to, acetyl, Fmoc, or Boc, and R' is an acid protecting group such as t-butyl or TSE. In certain preferred embodiments, the earbohydrate determinant is selected from the group consisting of Globo-H, fucosyl GM1, KH-1, glycophorin, STN, (2,3)ST, Le^y, N3, Tn, 2,6-STn, and TF. In certain other preferred embodiments, the carbohydrate determinant of the compound contains a Globo-H determinant or a fucosyl GM1 determinant, as described herein, as all or part of the carbohydrate determinant A.

In general, in preferred embodiments, the step of reacting an n-alkenyl glycoside under suitable conditions to generate an enamide ester comprises reacting an n-alkenyl glycoside first under oxidative conditions and second under olefination conditions in the presence of base (e.g., tetramethylguanidine) and phosphonate to generate an enamide ester.

Additionally, the step of reacting said enamide ester under suitable conditions to generate a glycoamino acid comprises reacting said enamide ester under hydrogenation conditions.

ĺ

5

10

15

20

25

In another aspect of the present invention, multi-antigenic glycopeptides are provided comprising a peptidic backbone made up of at least three glycoamino acids, wherein one or more of said amino acids are substituted with an n-alkyl glycosidic moiety having the structure:

wherein each occurrence of A is independently a carbohydrate determinant having the structure:

$$\begin{array}{c|c}
R_0 & & \\
\hline
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\$$

wherein a, b, c, d, e, f, g, h, i, x, y and z are independently 0, 1, 2 or 3, with the proviso that x, y and z are not simultaneously 0; wherein R₀ is hydrogen, a linear or branched chain alkyl, acyl, arylalkyl or aryl group; wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈ and R₉ are each independently hydrogen, OH, OR¹, NH₂, NHCOR¹, F, CH₂OH, CH₂OR¹, a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein R¹ is hydrogen, CHO, COOR¹¹, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group or a saccharide moiety having the structure:

$$\begin{array}{c|c}
R_0 & \xrightarrow{Z} & \xrightarrow{I_1} & \xrightarrow{I_2} & \xrightarrow{I_3} & \xrightarrow{I_4} & \xrightarrow{I_{13}} & \xrightarrow{I_{14}} & \xrightarrow{I_{15}} & \xrightarrow{I_{15$$

10

15

20

25

30

(

wherein Y and Z are independently NH or O; wherein k, l, r, s, t, u, v and w are each independently 0, 1 or 2; wherein R₁₀, R₁₁, R₁₂, R₁₃, R₁₄ and R₁₅ are each independently hydrogen, OH, ORⁱⁱⁱ, NH₂, NHCORⁱⁱⁱ, F, CH₂OH, CH₂ORⁱⁱⁱ, or a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein R₁₆ is hydrogen, COOH, COORⁱⁱ, CONHRⁱⁱ, a substituted or unsubstituted linear or branched chain alkyl or aryl group; wherein Rⁱⁱⁱ is hydrogen, CHO, COOR^{iv}, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group; and wherein Rⁱⁱ and R^{iv} are each independently H, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group;

wherein each occurrence of n is independently 0-8, whereby, if for each occurrence of n, n=0, at least one occurrence of A has a different structure from other occurrences of A; and wherein the n-alkyl glycosidic moiety is either α - or β -linked to an amino acid residue of the backbone. It will be appreciated that these inventive clustered glycopeptides are not limited to n-alkyl where n is greater than or equal to 1; rather multi-antigenic clustered glycopeptides can be linked via the traditional direct linkage (n=0) or via n-alkyl (such as pentyl) or any combination thereof. In other embodiments, each occurrence of A may be the same, however, n-alkenyl (n greater than 1) linkages are then utilized. In preferred embodiments, each occurrence of A is independently selected from the group consisting of Globo-H, fucosyl GM1, KH-1, glycophorin, STN, (2,3)ST, Le^y, N3, Tn, 2,6-STn, and TF.

In certain embodiments, trimeric antigenic clusters are desirable and thus the present invention also provides constructs attached to a linker via a carrier protein having the following structure:

(

wherein the linker is either a free carboxylic acid, (carboxamido)alkyl carboxamide, carboxamide, mono- or dialkyl carboxamide, MBS, diarylcarboxamide, linear or branched chain (carboxy)alkyl carboxamide, linear or branched chain (alkoxycarbonyl)alkyl-carboxamide, linear or branched chain branched (carboxy)arylalkylcarboxamide, chain (alkoxycarbonyl)alkylcarboxamide, an oligoester fragment comprising from 2 to about 20 hydroxy acyl residues, a peptidic fragment comprising from 2 to about 20 amino acyl residues, or a linear or branched chain alkyl or aryl carboxylic ester; wherein the carrier is a protein or lipid; wherein m is 1, 2 or 3; wherein RA, RB and RC are each independently H or methyl; and wherein RD, RE and RF are each independently an alkyl glycosidic moiety having the structure:

20

wherein each occurrence of A is independently selected from a carbohydrate domain having the structure:

10

15

20

(

$$\begin{array}{c|c}
R_0 & O & P_0 \\
\hline
 & P_$$

wherein a, b, c, d, e, f, g, h, i, x, y and z are independently 0, 1, 2 or 3, with the proviso that x, y and z are not simultaneously 0; wherein the carbohydrate domain is linked to the respective amino acyl or hydroxy acyl residue by substitution of a side group substituent selected from the group consisting of OH, COOH and NH₂; wherein R₀ is hydrogen, a linear or branched chain alkyl, acyl, arylalkyl or aryl group; wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈ and R₉ are each independently hydrogen, OH, OR¹, NH₂, NHCOR¹, F, CH₂OH, CH₂OR¹, a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein R¹ is hydrogen, CHO, COOR¹¹, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group or a saccharide moiety having the structure:

wherein Y and Z are independently NH or O; wherein k, l, r, s, t, u, v and w are each independently 0, 1 or 2; wherein R₁₀, R₁₁, R₁₂, R₁₃, R₁₄ and R₁₅ are each independently hydrogen, OH, ORⁱⁱⁱ, NH₂, NHCORⁱⁱⁱ, F, CH₂OH, CH₂ORⁱⁱⁱ, or a

substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein R₁₆ is hydrogen, COOH, COORⁱⁱ, CONHRⁱⁱ, a substituted or unsubstituted linear or branched chain alkyl or aryl group; wherein Rⁱⁱⁱ is hydrogen, CHO, COOR^{iv}, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group; and wherein Rⁱⁱ and R^{iv} are each independently H, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group; and wherein each occurrence of n is independently 0-8, whereby, if for each occurrence of n, n = 0, at least one occurrence of A has a different structure from other occurrences of A; and wherein the n-alkyl glycosidic moiety is either α- or β-linked to an amino acid.

In preferred embodiments, each occurrence of A is selected from the group consisting of Globo-H, fucosyl GM1, KH-1, glycophorin, STN, (2,3)ST, Le^y, N3, Tn, 2,6 STn and TF. In but one preferred example, the present invention provides a novel trimeric antigenic glycopeptide incorporating globo-H, Le^y and Tn, to generate a novel trimeric antigenic compound.

As detailed herein, in another aspect of the present invention, any of the inventive compounds may be conjugated to generate a glycoconjugate, and may be administered alone or with an immunological adjuvant for the treatment of the recurrence of cancer or may be administered alone or with an immunological adjuvant to induce antibodies in a subject.

25

35

10

15

20

Definitions

Certain compounds of the present invention, and definitions of specific functional groups are also described in more detail below. For purposes of this invention, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, Handbook of Chemistry and Physics, 75th Ed., inside cover, and specific functional groups are defined as described therein. Additionally, general principles of organic chemistry, as well as specific functional moieties and reactivity, are described in "Organic Chemistry", Thomas Sorrell, University Science Books, Sausalito: 1999, the entire contents of which are incorporated herein by reference.

It will be appreciated that the compounds, as described herein, may be substituted with any number of substitutents or functional moieties. In general, the term "substituted"

ί..

5

10

15

20

25

30

35

whether preceded by the term "optionally" or not, and substituents contained in formulas of this invention, refer to the replacement of hydrogen radicals in a given structure with the radical of a specified substituent. When more than one position in any given structure may be substituted with more than one substituent selected from a specified group, the substituent may be either the same or different at every position. As used herein, the term "substituted" is contemplated to include all permissible substituents of organic compounds. In a broad aspect, the permissible substituents include acyclic and cyclic, branched and unbranched, carbocyclic and heterocyclic, aromatic and nonaromatic substituents of organic compounds. For purposes of this invention, heteroatoms such as nitrogen may have hydrogen substituents and/or any permissible substituents of organic compounds described herein which satisfy the valencies of the heteroatoms. Furthermore, this invention is not intended to be limited in any manner by the permissible Combinations of substituents and variables substituents of organic compounds. envisioned by this invention are preferably those that result in the formation of stable compounds useful in the treatment of cancer, or in the inducement of antibodies, as described herein. The term "stable", as used herein, preferably refers to compounds which possess stability sufficient to allow manufacture and which maintain the integrity of the compound for a sufficient period of time to be useful for the purposes detailed herein.

The term "aliphatic", as used herein, includes both saturated and unsaturated, straight chain (i.e., unbranched), branched, cyclic, or polycyclic aliphatic hydrocarbons, which are optionally substituted with one or more functional groups. As will be appreciated by one of ordinary skill in the art, "aliphatic" is intended herein to include, but is not limited to, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, and cycloalkynyl moieties. Thus, as used herein, the term "alkyl" includes both straight, branched and cyclic alkyl groups. An analogous convention applies to other generic terms such as "alkenyl", "alkynyl" and the like. Furthermore, as used herein, the terms "alkyl", "alkynyl" and the like encompass both substituted and unsubstituted groups.

The term "alkylamino" refers to a group having the structure -NHR' wherein R' is alkyl, as defined herein. Examples of alkylamino include, but are not limited to, methylamino, ethylamino, iso-propylamino and the like.

5

10

15

20

25

30

Some examples of substituents of the above-described aliphatic (and other) moieties of compounds of the invention include, but are not limited to: F, Cl, Br, I, OH, NO₂, CN, C(O)-C₁-C₆-alkyl, C(O)-aryl, C(O)-heteroaryl, CO₂-alkyl, CO₂-aryl, CO₂-heteroaryl, CONH-2, CONH-C₁-C₆-alkyl, CONH-aryl, CONH-heteroaryl, OC(O)-C₁-C₆-alkyl, OC(O)-aryl, OC(O)-heteroaryl, OCO₂-alkyl, OCO₂-aryl, OCO₂-heteroaryl, OCONH-2, OCONH-C₁-C₆-alkyl, OCONH-aryl, OCONH-heteroaryl, NHC(O)-C₁-C₆-alkyl, NHC(O)-aryl, NHC(O)-heteroaryl, NHCO₂-alkyl, NHCO₂-aryl, NHCONH-heteroaryl, SO₂-C₁-C₆-alkyl, SO₂-aryl, C₃-C₆-cycloalkyl, CF₃, CH₂CF₃, CHCl₂, CH₂OH, CH₂CH₂OH, CH₂NH₂, CH₂SO₂CH₃, aryl, heteroaryl, benzyl, benzyloxy, aryloxy, heteroaryloxy, C₁-C₆-alkoxy, methoxymethoxy, methoxyethoxy, amino, benzylamino, arylamino, heteroarylamino, C₁-C₃-alkyl-amino, thio, aryl-thio, heteroarylthio, benzylthio, C₁-C₆-alkyl-thio, or methylthiomethyl. Additional examples of generally applicable substituents are illustrated by the specific embodiments shown in the Examples which are described herein.

In general, the terms "aryl" and "heteroaryl", as used herein, refer to stable monoor polycyclic, heterocyclic, polycyclic, and polyheterocyclic unsaturated moieties having preferably 3-14 carbon atoms, each of which may be substituted or unsubstituted. Substituents include, but are not limited to, any of the previously mentioned substitutents. i.e., the substituents recited for aliphatic moieties, or for other moieties as disclosed herein, resulting in the formation of a stable compound. In certain embodiments of the present invention, "aryl" refers to a mono- or bicyclic carbocyclic ring system having one or two aromatic rings including, but not limited to, phenyl, naphthyl, tetrahydronaphthyl, indanyl, indenyl and the like. In certain embodiments of the present invention, the term "heteroaryl", as used herein, refers to a cyclic aromatic radical having from five to ten ring atoms of which one ring atom is selected from S, O and N; zero, one or two ring atoms are additional heteroatoms independently selected from S, O and N; and the remaining ring atoms are carbon, the radical being joined to the rest of the molecule via any of the ring atoms, such as, for example, pyridyl, pyrazinyl, pyrimidinyl, pyrrolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, isooxazolyl, thiadiazolyl,oxadiazolyl, thiophenyl, furanyl, quinolinyl, isoquinolinyl, and the like.

5

20

25

It will be appreciated that aryl and heteroaryl groups (including bicyclic aryl groups) can be unsubstituted or substituted, wherein substitution includes replacement of one, two or three of the hydrogen atoms thereon independently with any one or more of the following moieties including, but not limited to: F, Cl, Br, I, OH, NO2, CN, C(O)-C1-C6-alkyl, C(O)-aryl, C(O)-heteroaryl, CO2-alkyl, CO2-aryl, CO2-heteroaryl, CONH2, CONH-C₁-C₆-alkyl, CONH-aryl, CONH-heteroaryl, OC(0)-C₁-C₆-alkyl, OC(0)-aryl, OC(O)-heteroaryl, OCO2-alkyl, OCO2-aryl, OCO2-heteroaryl, OCONH-C1-C6-alkyl, OCONH-aryl, OCONH-heteroaryl, NHC(0)-C1-C6-alkyl, NHC(0)-aryl, NHC(0)-heteroaryl, NHCO2-alkyl, NHCO2-aryl, NHCONH-heteroaryl, SO2-C1-C6-alkyl, SO₂-aryl, C₃-C₆-cycloalkyl, CF₃, CH₂CF₃, CHCl₂, CH₂OH, CH₂CH₂OH, CH₂NH₂, CH₂SO₂CH₃, aryl, heteroaryl, benzyl, benzyloxy, aryloxy, heteroaryloxy, C₁-C₆-alkoxy, methoxymethoxy, methoxyethoxy, amino, benzylamino, arylamino, heteroarylamino, C₁-C3-alkyl-amino, thio, aryl-thio, heteroarylthio, benzyl-thio, C1-C6-alkyl-thio, or Additional examples of generally applicable substitutents are illustrated by the specific embodiments shown in the Examples which are described herein.

The term "cycloalkyl", as used herein, refers specifically to groups having three to seven, preferably three to ten carbon atoms. Suitable cycloalkyls include, but are not limited to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and the like, which, as in the case of other aliphatic, heteroaliphatic or hetercyclic moieties, may optionally be substituted. F, Cl, Br, I, OH, NO₂, CN, C(O)-C₁-C₆-alkyl, C(O)-aryl, C(O)-heteroaryl, CO₂-alkyl, CO₂-aryl, CO₂-heteroaryl, CONH₂, CONH₂, CONH-C₁-C₆-alkyl, CONH-aryl, OCO₂-alkyl, OCO₂-aryl, OCO₂-heteroaryl, OCO)-to-C₆-alkyl, OC(O)-aryl, OC(O)-heteroaryl, OCO)-heteroaryl, OCONH-heteroaryl, NHC(O)-C₁-C₆-alkyl, NHC(O)-aryl, NHC(O)-heteroaryl, NHCO₂-alkyl, NHCO₂-alkyl, NHCO₂-aryl, NHCONH-heteroaryl, SO₂-C₁-C₆-alkyl, SO₂-aryl, C₃-C₆-cycloalkyl, CF₃, CH₂CF₃, CHCl₂, CH₂OH, CH₂CH₂OH, CH₂NH₂, CH₂SO₂CH₃, aryl, heteroaryl, benzyl, benzyloxy, aryloxy, heteroaryloxy, C₁-C₆-alkoxy, methoxymethoxy, methoxyethoxy, amino, benzylamino, arylamino, heteroarylamino, C₁-C₃-alkyl-amino, thio, aryl-thio, heteroarylthio, benzyl-thio, C₁-C₆-alkyl-thio, or methylthiomethyl. Additional examples

of generally applicable substitutents are illustrated by the specific embodiments shown in the Examples which are described herein.

(

10

15

25

30

The terms "halo" and "halogen" as used herein refer to an atom selected from fluorine, chlorine, bromine and iodine.

It will be appreciated that additional examples of generally applicable substitutents are illustrated by the specific embodiments shown in the Examples which are described herein, but are not limited to these Examples.

Description of the Drawing

Figure 1 depicts Fucosyl GM1, derivatives and constructs thereof.

Figure 2 depicts synthesis of the ABC trisaccharide 4 and depicts the thioethyl donor 5.

Figure 3 depicts the synthesis of hexasaccharide 6a and the synthesis of Fucosyl GM₁ pentenyl glycoside 1b. Reagents: (a) MeOTf, CH₂Cl₂:Et₂O (2:1), 0°C, 23%; (b) (i) DMDO, CH₂Cl₂; (ii) PnOH, ZnCl₂, -78°C, 65%; (c) TBAF, AcOH, THF; (d) NaOMe, MeOH; (e) NaOH, THF; (f) Na/NH₃, THF -78°C, then MeOH; (g) Ac₂O, pyridine, DMAP, CH₂Cl₂, 46% 5 steps.

Figure 4 depicts the synthesis of trisaccharide acceptor 15. Reagents: (a) Ag₂CO₃, cat. I₂, PnOH, CH₂Cl₂, 75%; (b) NaOMe, MeOH; (c) Acetone, cat. PPTS, 44% 2 steps; (d) BnBr, NaH, DMF; 84%; (e) 80% AcOH: H₂O, 90%; (f) 3, TMSOTf, EtCN, molecular sieves, -40°C, 77%.

Figure 5 depicts the synthesis of Fucosyl GM₁ Pentenyl Glycoside. Reagents: (a) MeOTf, CH₂Cl₂:Et₂O, 0°C, 70%; (b) TBAF, AcOH, THF; (c) NaOMe, MeOH; (d) NaOH, THF; (e) Na/NH₃, THF, -78°C, then MeOH; (f) Ac₂O, pyridine, DMAP, CH₂Cl₂, 45% 5 steps, (g) steps c-d, 96%.

Figure 6 depicts the synthesis of Fucosyl GM₁ KLH conjugate 1c.

Figure 7 depicts Globo-H, derivatives and constructs thereof.

Figure 8 depicts a synthetic scheme for a second generation synthesis of globo-H and constructs thereof.

Figure 9 depicts a retrosynthetic analysis of Globo-H and conjugates thereof.

(

5

15

20

25

30

35

Figure 10 depicts the synthesis of glycoside 25 and thioethyl donor 28. Reagents: (a) HBr, Ac₂O, AcOH, 96%; (b) PentOH, Ag₂CO₃, CH₂Cl₂, 4Å molecular sieves, 75%; c) NaOMe, MeOH; then Dowex-H⁺; (d) BnBr, Bu₂SnO, Bu₄NI, C₆H₆, 54% two steps; e) PhCH(OMe)₂, CSA, CH₃CN, 72%; (f) BnBr, NaH, DMF, Et₄NI, 97%; (g) NaCNBH₃, HCl, Et₂O, THF, 79%; (h) DMDO, CH₂Cl₂; (i) HF/pyridine, 85% two steps; (j) BnBr, NaH, DMF, 95%; (k) Cp₂Zr(OTf)₂, toluene/THF 5:1, 80% (α), α:β 10:1; (l) DDQ, CH₃CN, H2O, 84%.

Figure 11 depicts the synthesis of Globo-H Pentenyl Glycoside (16c).

Figure 12 depicts the conjugation of Globo-H to carrier protein KLH.

Figure 13 depicts immunoconjugation of tumor antigens Globo-H and Fucosyl GM₁ and the developed glycoamino acid sequence.

Figure 14 depicts the synthesis of peracetylated lactose amino acid derivatives.

Figure 15 depicts the synthesis of a peptide containing the Tn antigen, Lewis^y antigen, and the MBr1 antigen. Reagents: (a) TBAF, THF; (b) AcSCH₂C(O)(CH₂)₃NH₂, BOP reagent, iPr₂NEt, 54%, 2 steps; (c) TFA, CH₂Cl₂; (d) BOP reagent, iPr₂NEt, 86%, 2 steps; (e) 52, BOP reagent, iPr₂NEt, 64%, 2 steps; (f) Ac₂O, Et₃N, cat. DMAP, 95%, 2 steps.

Figure 16 depicts the preparation of fully deprotected glycopeptide 54. Figure 17 depicts the synthesis of α -Tn pentenyl glycoside 40.

Detailed Description of the Invention

As discussed above, the desire to develop improved methods for the preparation of fully synthetic vaccines has led to increased research efforts directed toward the synthesis of naturally occurring complex carbohydrate antigens, as well as novel complex structures (e.g., glycopeptides) incorporating these antigenic structures. As is often the case during the course of any such large synthetic undertaking, improved synthetic methods are often developed that can be applied universally. In particular, synthetic studies of naturally occurring antigenic structures has led to the development of novel methodologies enabling the development of heretofore unavailable synthetic carbohydrate-based vaccines. For a review, see Danishefsky, S.J.; Allen, J.R., Angew. Chem. Int. Ed. Engl. 2000, 39, 836-863, and references cited therein.

(

5

10

15

20

25

Significantly, the present invention provides improved methodologies for the synthesis of complex carbohydrates and related therapeutic compounds (e.g., glycoconjugates and/or glycopeptides). In particular, in the context of synthetic studies developed for the total syntheses of fucosyl GM1 and the n-pentenyl glycoside of Globo-H, generalized methodologies were developed for the improved synthesis of complex carbohydrate structures. This general synthetic method involves the realization that the incorporation of an n-alkenyl glycoside protecting group at the reducing end of a carbohydrate acceptor allows for increased coupling efficiencies and accessibility to complex carbohydrates. In yet another aspect, the present invention also provides the recognition that for certain protected carbohydrates, the n-alkenyl moieties can serve as useful precursors that can be utilized ultimately for the synthesis of complex glycopeptides.

Furthermore, the present invention also provides the recognition that the presence of the n-alkenyl moiety, whether or not in the context of an antigenic n-pentenyl glycoside or glycopeptide, is advantageous for the development of improved carbohydrate based therapeutics (e.g., fully synthetic vaccines) because more efficient syntheses of conjugation precursors can be prepared (and ultimately conjugated), and the n-alkenyl carbohydrate also serves as a precursor for the synthesis of novel n-alkyl glycoamino acids, as described herein. The ability to easily access these glycoamino acids allows for the ultimate synthesis of complex clustered glycopeptides. Significantly, the methodologies provided by the present invention, as described above and in more detail herein, allow the efficient preparation of complex glycopeptide structures having more than one type of carbohydrate determinant.

Specific examples, particularly with respect to the total synthesis of fucosyl GM1 and a novel synthetic scheme for the synthesis of the n-pentenyl glycoside of Globo-H are described in more detail below, along with certain general methodologies developed during the course of these syntheses. It will be appreciated by one of ordinary skill in the art that these examples are not intended to be limiting; rather all equivalents are intended to be incorporated into the scope of the present invention.

20

25

35

5 Inventive Compounds and Methods for the Synthesis Thereof

As mentioned, the total synthesis of complex antigenic structures has led to significant development in methodologies for complex carbohydrate synthesis. Of particular recent interest is the naturally occurring antigenic structure, fucosylated GM1 ganglioside as shown in Figure 1 (1a) which heretofore had not yet been synthesized. Nilsson et al. identified fucosyl GM1 as a specific marker associated with small lung cancer (SCLC) cells (Nilsson et al., Glycoconjugate J. 1984, 1, 43; Brezicka et al., Cancer Res. 1989, 49, 1300). These workers isolated the glycosphingolipid fucosyl GM1 (1a) as the major ganglioside component contained in human SCLC tissue. Furthermore, monoclonal antibodies (F12) to the antigen serve to detect fucosyl GM1 in tissues and serum of SCLC patients (Nilsson et al., Cancer Res. 1986, 46, 1403; Vangsted et al., Cancer Res. 1991, 51, 2897). Immunohistochemistry studies have suggested that, due to its highly restricted distribution in normal tissues, fucosyl GM1 could be an excellent target for immune attack against SCLC. Remarkably, fucosyl GM1 has thus far not been found on any other human cancer cell lines, indicating that it is very SCLC tumor specific (Zhang et al., Int. J. Cancer 1997, 73, 42).

The structural assignment of the carbohydrate moiety of the SCLC antigen was based on a combination of enzymatic and chemical degradations (Nilsson et al., Glycoconjugate J. 1984, 1, 43). While there was no particular reason to question this assignment, the development of a carbohydrate based attack on SCLC could benefit from a definitive assignment of the linkage modes of the various monosaccharides, including the stereochemistry at each glycosidic attachment. Furthermore, no syntheses of this carbohydrate sector have appeared in the literature. In a preferred embodiment, a synthetic scheme would allow for presentation of the hexasaccharide epitope independent of the ceramide to the F12 mAb to ensure that all specificity is directed at the carbohydrate sector. In other preferred embodiments, the construct should be so functionalized as to anticipate the need for its conjugation to a carrier protein in anticipation of building an effective anti-tumor vaccine. As detailed herein, the ability to generate an n-alkenyl glycoside enables for the efficient synthesis of this epitope and allows for its effective modification and/or conjugation to build an effective anti-tumor vaccine.

Thus, in one aspect of the present invention, the synthesis of the complex fucosyl GM1 carbohydrate sector has been achieved and a compound having the structure as shown below is provided:

10

15

5

wherein each occurrence of R' is independently hydrogen or a protecting group; wherein each occurrence of R" is independently hydrogen or a nitrogen protecting group; wherein R is hydrogen, substituted or unsubstituted alkyl, alkenyl, -NHR", wherein R" is a protein, peptide or lipid linked to N directly or through a crosslinker, an amino acyl moiety, an amino acyl residue of a peptide, an amino acyl residue of a protein, which amino acyl moiety or residue, or -NHR" is linked to O via a polymethylene chain having the structure -(CH₂)_r, where r is an integer between 1 and 9, or wherein R is substituted with a moiety having the structure:

20

In certain preferred embodiments, each occurrence of R' is hydrogen. In certain other preferred embodiments of the present invention, R is n-alkenyl, including, but not

limited to allyl, propenyl, butenyl and pentenyl. In a particularly preferred embodiment, R is n-pentenyl. In certain other preferred embodiments, R is -NHR'", an amino acyl moiety, an amino acyl residue of a peptide, or an amino acyl residue of a protein, as described above, wherein r is preferably 4. In still other preferred embodiments, a compound as described above is provided, with the proviso that the compound is not the glycosphingolipid structure.

In another aspect of the present invention, a method for the synthesis of fucosyl GM1 glycoside is provided, said method comprising the steps of:

(a) providing a thioethyl donor having the structure:

15

wherein P is a protecting group;

(b) providing a trisaccharide acceptor having the structure:

20

wherein n is 0-8, and wherein P is a protecting group; and

5

10

15

20

(c) reacting said thioethyl donor and said trisaccharide acceptor under conditions to generate a protected hexasaccharide and subsequently deprotecting the protected hexasaccharide under suitable conditions to generate n-alkenyl fucosyl GM1 glycoside.

In yet another aspect of the present invention, novel derivatives of Globo-H are provided and a novel general synthetic methodology for the synthesis thereof. The derivatives of Globo-H are depicted below:

wherein each occurrence of R' is hydrogen or a protecting group, wherein R" is hydrogen or a nitrogen protecting group; wherein R is hydrogen, substituted or unsubstituted alkyl or alkenyl, wherein the alkenyl moiety has four or more carbons; -NHR", wherein R'" is a protein, peptide or lipid linked to N directly or through a crosslinker; an amino acyl moiety; an amino acyl residue of a peptide; an amino acyl residue of a protein; which amino acyl moiety or residue or -NHR" is linked to O via a polymethylene chain having the structure -(CH₂)_r, where, if said carbohydrate moiety is linked to O via an α -linkage, r is an integer between 2 and 9, or, alternatively, if said carbohydrate moiety is linked to O via a β -linkage, r is an integer between 1 and 9; or wherein R is substituted with a moiety having the structure:

25

10

15

In certain preferred embodiments, each occurrence of R' is hydrogen. In certain other preferred embodiments of the present invention, R is n-alkenyl, including, but not limited to allyl, propenyl, butenyl and pentenyl. In a particularly preferred embodiment, R is n-pentenyl. In certain other preferred embodiments, R is an amino acyl moiety, an amino acyl residue of a peptide, or an amino acyl residue of a protein, as described above, wherein r is preferably 4. In still other preferred embodiments, a compound as described above is provided, with the proviso that the compound is not the glycosphingolipid structure.

As described in more detail herein in Example 2, a similar methodology to that described for fucosyl GM1 is employed for the synthesis of Globo-H and derivatives thereof. Thus, in another aspect of the present invention, a method for the improved synthesis of Globo-H, and derivatives thereof, said method comprising the steps of:

(a) providing a thioethyl donor having the structure:

20

wherein P is a protecting group; and

(b) providing a trisaccharide acceptor having the structure:

(

5

10

15

20

25

30

wherein n is 0-8, and wherein P is a suitable protecting group; and

(c) reacting said thioethyl donor and said trisaccharide acceptor under conditions to generate a protected hexasaccharide and subsequently deprotecting the protected hexasaccharide under suitable conditions to generate n-alkenyl Globo-H.

It will be appreciated that for each of the methods as detailed herein, the full arsenal of protecting groups known in the art of organic synthesis can be utilized, for example, as set forth in "Activating Agents and Protecting Groups: Handbook of Reagents for Organic Synthesis" Roush, W.R. and Pearson, A.J., Eds., John Wiley & Sons: 1999; and "Protective Groups in Organic Synthesis" Greene, T.W. and Wuts, P.G., John Wiley & Sons, New York: 1999, the entire contents of which are hereby incorporated by reference. In but a few examples, suitable protecting groups utilized herein include, but are not limited to, Bn (benzyl), TIPS (triisopropylsilyl), and Ac (acetate). In a preferred embodiment of the present invention, said thioethyl donor and said trisaccharide acceptor are reacted under MeOTf promotion, as described herein. It will be appreciated by one of ordinary skill in the art however, that a variety of conditions known in the art of organic synthesis can be utilized to effect coupling of these moieties.

It will also be appreciated that the novel n-alkenyl moieties provided herein can be subsequently modified to generate useful compounds (e.g., alkyl derivatives and glycoamino acids) or constructs thereof (e.g., glycopeptides and conjugated derivatives).

In addition to providing the first synthesis of fucosyl GM1 and improved synthetic methodologies for Globo-H, as described above, in a more general aspect, the present invention provides novel synthetic methodologies for the synthesis of complex carbohydrates comprising (1) providing a carbohydrate acceptor having a reducing end alkenyl group; (2) providing a suitable donor compound and (3) coupling said donor and

acceptor under conditions to generate an alkenyl glycoside. Using this method, complex antigenic alkenyl glycosides are provided, as described above, many of which never before have been provided, which can then be conjugated or further reacted, as described herein, to generate glycoconjugates and glycopeptide structures.

Thus, in general, the present invention provides novel compounds and/or conjugates having the general structure:

wherein R is hydrogen; substituted or unsubstituted alkyl; alkenyl; aryl;
-CH₂CH(CO₂R')(NHR"), wherein R' or R" are each indpendently hydrogen, protecting
group, substituted or unsubstituted alkyl, a linker, aryl, peptide, protein or lipid; or
NHR", wherein R" is a protein, peptide, or lipid, linked to N directly or through a
crosslinker; wherein n is 0-8; wherein A is a carbohydrate domain having the structure:

$$\begin{array}{c|c}
R_0 & & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_4 & \\
\hline
 & R_6 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_4 & \\
\hline
 & R_5 & \\
\hline
 & R_6 & \\
\hline
 & R_6 & \\
\hline
 & R_7 & \\
\hline
 & R_8 &$$

20

25

15

wherein a, b, c, d, e, f, g, h, i, x, y and z are independently 0, 1, 2 or 3, with the proviso that x, y and z are not simultaneously 0; wherein R_0 is hydrogen, a linear or branched chain alkyl, acyl, arylalkyl or aryl group; wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 and R_9 are each independently hydrogen, OH, OR¹, NH₂, NHCOR¹, F, CH₂OH, CH₂OR¹, a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein R^1 is

15

20

25

30

hydrogen, CHO, COORⁱⁱ, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group or a saccharide moiety having the structure:

wherein Y and Z are independently NH or O; wherein k, l, r, s, t, u, v and w are each independently 0, 1 or 2; wherein R₁₀, R₁₁, R₁₂, R₁₃, R₁₄ and R₁₅ are each independently hydrogen, OH, ORⁱⁱⁱ, NH₂, NHCORⁱⁱⁱ, F, CH₂OH, CH₂ORⁱⁱⁱ, or a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein R₁₆ is hydrogen, COOH, COORⁱⁱ, CONHRⁱⁱ, a substituted or unsubstituted linear or branched chain alkyl or aryl group; wherein Rⁱⁱⁱ is hydrogen, CHO, COOR^{iv}, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group; and wherein Rⁱⁱ and R^{iv} are each independently H, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group;

with the proviso that if A is KH-1, N3, globo-H, glycophorin, Tn, TF, STN, (2,3-ST), 2,6-STn or Le^y, and A is α -O-linked, then n is at least 1.

In certain preferred embodiments of the present invention, R is allyl, n is 2 and thus the inventive compound is a n-pentenyl moiety. In certain other embodiments of the present invention, R is NHR'", and the protein R" is KLH or Bovine Serine Albumin. In still other embodiments of the present invention, R is NHR'", and the lipid R" is PamCys. It will be appreciated that the protein or lipid can be linked to N directly or through a crosslinker and thus R" incorporates proteins, peptides and lipids, as well as (crosslinker-protein), (crosslinker-peptide) and (crosslinker-lipid) moieties. In certain preferred embodiments, the crosslinker is MMCCH (4-(maleimidomethyl) cyclohexane-1-carboxyl hydrazide).

In still other embodiments, the inventive compound is a glycoamino acid and thus R is CH₂CH(CO₂R')(NHR"), which compound has the structure:

10

15

20

25

30

5

In certain preferred embodiments, the glycoamino acids of the present invention are derived from n-pentenyl glycosides and thus n is 3. In certain other preferred embodiments, R' and R" are a protecting group, each independently selected from the group consisting of t-butyl, TSE, Boc, Fmoc and acetyl.

For each of the compounds described above, in certain preferred embodiments the carbohydrate determinant is selected from the group consisting of Globo-H, fucosyl GM1, KH-1, N3, glycophorin, Tn, TF, STN, (2,3)ST, 2,6-STn, and Le^y. In other preferred embodiments, the carbohydrate determinant of the compound contains a Globo-H determinant or fucosyl GM1 determinant as all of part of the carbohydrate determinant A.

As described above, specifically in the context of the second generation synthesis of the MBr1 antigen (GloboH) and the total synthesis of the fucosylated ganglioside of GM1 (fucosyl GM1), incorporating the reducing end n-alkenyl moiety (specifically n-pentenyl) offers certain benefits. First, the anomeric n-pentenyl glycoside linkage serves as an effective linker for immunoconjugation to carrier protein KLH and also provides some advantages in terms of synthetic convergency. In the context of protected carbohydrates, the n-alkenyl moieties are also capable of acting as donors for glycosylation (see, for example, Fraser –Reid et al., SynLett, 1992, 927).

In this context, the present invention additionally provides methods for the synthesis of n-alkyl glycoamino acids, as described in more detail below for Globo-H and fucosyl GM1 and their subsequent use to generate glycopeptides and synthetic constructs thereof.

In general, the inventive method for the production of these glycoamino

subjecting said alkenyl glycosidic moiety to oxidative conditions to generate an aldehyde;

3) subjecting said aldehyde to olefination conditions to generate an enamide ester; 4) subjecting said resulting enamide ester to conditions sufficient to hydrogenate said enamide ester to generate a protected glycoamino acid and 5) deprotecting said protected glycoamino acid under suitable conditions to generate a desired glycoamino acid.

In particular, a novel method for the synthesis of a glycoamino acid, the structure of which is set forth herein, is provided, which comprises the steps of:

(a) providing an alkenyl glycoside having the structure:

15

(b) reacting said alkenyl glycoside under suitable conditions to generate an enamide ester having the structure:

20

(b) reacting said enamide ester under suitable conditions to generate a glycoamino acid having the structure:

25

wherein, for each of the structures above, n is 0-8, wherein A is a carbohydrate domain having the structure:

30

$$R_0$$
 R_0
 R_0

10

wherein a, b, c, d, e, f, g, h, i, x, y and z are independently 0, 1, 2 or 3, with the proviso that x, y and z are not simultaneously 0; wherein R₀ is hydrogen, a linear or branched chain alkyl, acyl, arylalkyl or aryl group; wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈ and R₉ are each independently hydrogen, OH, OR¹, NH₂, NHCOR¹, F, CH₂OH, CH₂OR¹, a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein R¹ is hydrogen, CHO, COOR¹¹, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group or a saccharide moiety having the structure:

15

... **20**

wherein Y and Z are independently NH or O; wherein k, l, r, s, t, u, v and w are each independently 0, 1 or 2; wherein R₁₀, R₁₁, R₁₂, R₁₃, R₁₄ and R₁₅ are each independently hydrogen, OH, ORⁱⁱⁱ, NH₂, NHCORⁱⁱⁱ, F, CH₂OH, CH₂ORⁱⁱⁱ, or a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein R₁₆ is hydrogen, COOH, COORⁱⁱ,

20

25

30

35

CONHRⁱⁱ, a substituted or unsubstituted linear or branched chain alkyl or aryl group; wherein Rⁱⁱⁱ is hydrogen, CHO, COOR^{iv}, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group; and wherein Rⁱⁱ and R^{iv} are each independently H, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group;

and wherein for the glycoamino acid structure R' and R" are each independently hydrogen, protecting group, substituted or unsubstituted alkyl, linker, aryl, peptide, protein or lipid; or NHR", wherein R" is a protein, peptide, or lipid, linked to N directly or through a crosslinker. In preferred embodiments, R' and R" are each independently hydrogen or a protecting group. In particularly preferred embodiments R" is a nitrogen protecting group, including, but not limited to, acetyl, Fmoc, Boc, and R' is an acid protecting group such as t-butyl or TSE. It will be appreciated, however, that a variety of protecting groups known in the art of organic synthesis can be employed, as referenced herein.

In certain preferred embodiments, the carbohydrate determinant is selected from the group consisting of Globo-H, fucosyl GM1, KH-1, glycophorin, STN, (2,3)ST, Le^y, N3, Tn, 2,6-STn, and TF. In other preferred embodiments, the carbohydrate determinant of the compound contains a Globo-H determinant or fucosyl GM1 determinant as all or part of the carbohydrate determinant A.

In general, in preferred embodiments, the step of reacting an n-alkenyl glycoside under suitable conditions to generate an enamide ester comprises reacting an n-alkenyl glycoside first under oxidative conditions and second under olefination conditions in the presence of base (e.g., tetramethylguanidine) and phosphonate to generate an enamide ester. It will be appreciated that other oxidative conditions known in the art of organic synthesis can be employed, including, but not limited to OsO₄ and periodate, or OsO₄ and Pb(OAc)₄. Additionally, other well-known bases can be utilized in the present invention, including, but not limited to, lithium t-butoxide or lithium hexamethyl disilylazide.

In preferred embodiments, reacting said enamide ester under suitable conditions to generate a glycoamino acid comprises reacting said enamide ester under hydrogenation conditions and subsequent reaction under deprotection conditions to generate a glycoamino acid. It is particularly preferred that the hydrogenation conditions employed

10

15

20

25

are asymmetric hydrogenation conditions. In a preferred embodiment, asymmetric hydrogenation can be achieved by utilizing an ethyl DuPHOS catalyst precursor, as described in more detail herein (see, Burk et al. Accts. Chem. Res. 2000, 33, 3631; Burk et al. Pure & Appl. Chem. 1996, 68, 37).

It will be appreciated that the ability to generate the glycoamino acids, as described herein, ultimately enables the synthesis of novel clustered glycopeptides, a motif commonly found on the surface of cancer cells (mucin-like structures) which are desirable for the uses described herein as anticancer vaccines. For example, immunological studies indicate that, in general, the clustering of antigens in glycopeptides results in a more therapeutically immune response than with singly glycosylated peptides (see, Lo-Man, R. et al., Cancer Res., 1999, 59, 1520; Reddish et al., Glycoconjugate J. 1997, 14, 549).

To date, the clustering of α -O-linked antigens has been accomplished with the same antigen across the peptide backbone via the traditional allyl linkage, as described in pending US patent numbers 09/083,776 and 09/276,595, the entire contents of which are hereby incorporated by reference. However, the present invention efficiently provides peptides having different antigens simultaneously in a clustered format. Thus, in one aspect, the present invention provides a multi-antigenic glycopeptide comprising a peptidic backbone made up of at least three glycoamino acids, wherein one or more of said amino acids are substituted with an n-alkyl glycosidic moiety having the structure:

wherein each occurrence of A is independently a carbohydrate determinant having the structure:

Ĭ.

5

10

15

20

$$R_0$$
 R_0 R_0

wherein a, b, c, d, e, f, g, h, i, x, y and z are independently 0, 1, 2 or 3, with the proviso that x, y and z are not simultaneously 0; wherein R₀ is hydrogen, a linear or branched chain alkyl, acyl, arylalkyl or aryl group; wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈ and R₉ are each independently hydrogen, OH, ORⁱ, NH₂, NHCORⁱ, F, CH₂OH, CH₂ORⁱ, a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein Rⁱ is hydrogen, CHO, COORⁱⁱ, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group or a saccharide moiety having the structure:

wherein Y and Z are independently NH or O; wherein k, l, r, s, t, u, v and w are each independently 0, 1 or 2; wherein R₁₀, R₁₁, R₁₂, R₁₃, R₁₄ and R₁₅ are each independently hydrogen, OH, ORⁱⁱⁱ, NH₂, NHCORⁱⁱⁱ, F, CH₂OH, CH₂ORⁱⁱⁱ, or a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein R₁₆ is hydrogen, COOH, COORⁱⁱ, CONHRⁱⁱ, a substituted or unsubstituted linear or branched

(

10

20

25

30

PCT/US00/22894

chain alkyl or aryl group; wherein Riii is hydrogen, CHO, COORiv, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group; and wherein Rii and Riv are each independently H, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group;

wherein each occurrence of n is independently 0-8, whereby, if for each occurrence of n, n = 0, at least one occurrence of A has a different structure from other occurrences of A; and wherein the n-alkyl glycosidic moiety is either α- or β-linked to an amino acid residue of the backbone. It will be appreciated that these inventive clustered glycopeptides are not limited to n-alkyl where n is greater than or equal to 1; rather multiantigenic clustered glycopeptides can be linked via the traditional direct linkage (n=0) or 15 via n-alkyl (such as pentyl) or any combination thereof. In preferred embodiments, each occurrence of A is independently selected from the group consisting of Globo-H, fucosyl GM1, KH-1, glycophorin, STN, (2,3)ST, Le^y, N3, Tn, 2,6-STn, and TF.

It will also be appreciated from the structure as set forth above, that, in addition to providing multi-antigenic structures, the present invention additionally provides clustered structures having n-alkyl linkages. Thus, in yet another aspect of the present invention, n-alkyl linked (where n is greater than or equal to 1) clustered glycopeptides are provided, which glycopeptides may incorporate multiple antigenic structures or may also incorporate all of the same antigenic structures.

In general, the generation of the inventive glycopeptides comprises treating a first glycoamino acid with a deprotecting agent to reveal the corresponding carboxylic acid and then coupling said carboxylic acid under suitable conditions with a spacer moiety and a protecting group to generate a protected amide. A second glycoamino acid can then be coupled under standard conditions (e.g., BOP promoter or other known coupling reagents known in the art of peptide couplings) these couplings can be continued until a peptide of desired length is obtained. It will also be appreciated that solid phase methods of peptide synthesis known in the art can also be employed in the method of the present invention to generate the inventive glycopeptides.

While the glycopeptide of the present invention is not intended to be limited in size, in certain preferred embodiments, trimeric antigenic clusters are desirable and thus

the present invention also provides constructs attached to a linker via a carrier protein having the following structure:

wherein the linker is either a free carboxylic acid, (carboxamido)alkyl carboxamide, 10 or dialkyl carboxamide, monoprimary carboxamide, monodiarylcarboxamide, linear or branched chain (carboxy)alkyl carboxamide, linear or branched chain (alkoxycarbonyl)alkyl-carboxamide, linear or branched chain (carboxy)arylalkylcarboxamide, branched chain linear or (alkoxycarbonyl)alkylcarboxamide, an oligoester fragment comprising from 2 to about 20 hydroxy acyl residues, a peptidic fragment comprising from 2 to about 20 amino acyl residues, or a linear or branched chain alkyl or aryl carboxylic ester; wherein the carrier is a protein or lipid; wherein m is 1, 2 or 3; wherein RA, RB and RC are each independently H or methyl; and wherein RD, RE and RF are each independently an alkyl glycosidic moiety having the structure: 20

wherein each occurrence of A is independently selected from a carbohydrate domain having the structure:

5

10

15

wherein a, b, c, d, e, f, g, h, i, x, y and z are independently 0, 1, 2 or 3, with the proviso that x, y and z are not simultaneously 0; wherein the carbohydrate domain is linked to the respective amino acyl or hydroxy acyl residue by substitution of a side group substituent selected from the group consisting of OH, COOH and NH₂; wherein R₀ is hydrogen, a linear or branched chain alkyl, acyl, arylalkyl or aryl group; wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈ and R₉ are each independently hydrogen, OH, OR¹, NH₂, NHCOR¹, F, CH₂OH, CH₂OR¹, a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein R¹ is hydrogen, CHO, COOR¹¹, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group or a saccharide moiety having the structure:

$$\begin{array}{c|c} R_{0} & \begin{array}{c} Z & \\ \end{array} & \begin{array}{c} \\ \end{array} & \end{array} & \begin{array}{c} \\ \end{array} & \end{array} & \begin{array}{c} \\ \end{array} & \begin{array}{c}$$

20

(:

wherein Y and Z are independently NH or O; wherein k, l, r, s, t, u, v and w are each independently 0, 1 or 2; wherein R_{10} , R_{11} , R_{12} , R_{13} , R_{14} and R_{15} are each independently hydrogen, OH, OR^{lii}, NH₂, NHCOR^{lii}, F, CH₂OH, CH₂OR^{lii}, or a

substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein R_{16} is hydrogen, COOH, COORⁱⁱ, CONHRⁱⁱ, a substituted or unsubstituted linear or branched chain alkyl or aryl group; wherein R^{iii} is hydrogen, CHO, COOR^{iv}, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group; and wherein R^{ii} and R^{iv} are each independently H, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group; and wherein each occurrence of n is independently 0-8, whereby, if for each occurrence of n, n = 0, at least one occurrence of A has a different structure from other occurrences of A; and wherein the n-alkyl glycosidic moiety is either α - or β -linked to an amino acid.

In certain embodiments, each occurrence of A is independently selected from the group consisting of Globo-H, fucosyl GM1, KH-1, glycophorin, STN, (2,3)-ST, Le^y, N3, Tn, 2,6-STn and TF. In but one preferred example, the present invention provides a novel trimeric antigenic glycopeptide incorporating globo-H, Le^y and Tn, to generate a novel trimeric antigenic compound, as described in more detail in Example 3 herein.

20

25

30

35

10

15

Pharmaceutical Compositions, Constructs and Uses Thereof

As described above, the present invention provides compounds and synthetic methodologies useful in the development of novel therapeutic agents, particularly for fully synthetic cancer vaccines. In general, the compounds and glycopeptides prepared as disclosed herein can be conjugated to a protein carrier or a lipid to generate useful glycoconjugates for the treatment and prevention, (preferably the prevention of the recurrence), of cancer in a subject suffering therefrom. In addition, glycoconjugates prepared by processes disclosed herein are antigens useful in adjuvant therapies as vaccines capable of inducing antibodies immunoreactive with various tumor cells. Such adjuvant therapies may reduce the rate of recurrence of certain cancers, and increase survival rates after surgery. Clinical trials on patients surgically treated for cancer who are then treated with vaccines prepared from a cell surface differentiation antigen found in patients lacking the antibody prior to immunization, a highly significant increase in disease-free interval may be observed. Cf. P.O. Livingston, et al., J. Clin. Oncol., 1994, 12, 1036.

Thus, the present invention provides pharmaceutical compositions for treating cancer, preferably for preventing the recurrence of cancer, comprising any of the compounds of the present invention disclosed herein, as an active ingredient, optionally, though typically in combination with a pharmaceutically acceptable carrier. The pharmaceutical compostions of the present invention may further comprise other therapeutically active ingredients.

5

10

15

25

35

This method of treatment comprises administering to the subject a therapeutically effective amount of any of the glyconjugates disclosed herein, optionally in combination with a pharmaceutically acceptable carrier. The method may be applied wherein the cancer is a solid tumor or an epithelial tumor. As mentioned above, methods for the treatment of cancer (preferably for the prevention of recurrence of cancer) are provided, as well as methods for inducing antibodies in a human subject, wherein the antibodies are capable of specifically binding with human tumor cells, which comprises administering to the subject an amount of any of the glycoconjugates disclosed above effective to induce antibodies. In certain embodiments, the carbohydrate antigen is linked to an effective carrier either directly or through a crosslinker, which carrier is a protein or lipid. In certain embodiments, the carrier protein is bovine serine albumin, polylysine or KLH. In certain other embodiments, the lipid is PamCys.

In addition, the present invention provides the related method of inducing antibodies which further comprises co-administering an immunological adjuvant, or a combination of immunological adjuvants. In certain embodiments, the adjuvant is a saponin adjuvant (see, e.g., Marciani et al., Vaccine, 2000, 18, 3141, US Patent No.: 6,080,725 and 5,977,081, the entire contents of which are hereby incorporated by reference). One example of a preferred saponin adjuvant includes, but is not limited to, GPI-0100, (Galenica Pharmaceuticals, Inc., Frederick, MD) which is a semi-synthetic adjuvant derived by modifying selected natural saponins. In certain other preferred embodiments, the adjuvant is bacteria or liposomes. In certain examples, the adjuvant includes but is not limited to, Salmonella minnesota cells, bacille Calmette-Guerin or QS21.

It will be appreciated that the magnitude of the therapeutic dose of the compounds of the invention will vary with the nature and severity of the condition to be treated and

with the particular compound and its route of administration. In general, the daily dose range for anticancer activity lies in the range of 0.0001 to 1.0 mg/kg of body weight in a mammal, although the present invention is not intended to be limited by this range.

(

10

15

20

25

30

35

Any suitable route of administration may be employed for providing a mammal, especially a human, with an effective dosage of a compound disclosed herein. For example, oral, rectal, topical, parenteral, ocular, pulmonary, nasal, etc. routes may be employed. Dosage forms include tablets, troches, dispersions, suspensions, solutions, capsules, creams, ointments, aerosols, etc. In preferred embodiments, the effective dosage is employed using a syringe injection.

The inventive compositions include those suitable for oral, rectal, topical (including transdermal devices, aerosols, creams, ointments, lotions and dusting powders), parenteral (including subcutaneous, intramuscular, and intravenous), ocular (opthalmic), pulmonary (nasal or buccal inhalation) or nasal administration. Although the most suitable route in any given case will depend largely on the nature and severity of the condition being treated and on the nature of the active ingredient. They may be conveniently presented in unit dosage form and prepared by any of the methods well known in the art of pharmacy.

In preparing oral dosage forms, any of the unusual pharmaceutical media may be used, such as water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents, and the like in the case of oral liquid preparations (e.g., suspensions, elixers and solutions); or carriers such as starches, sugars, microcrystalline cellulose, diluents, granulating agents, lubricants, binders, disinterating agents, etc., in the case of oral solid preparations are preferred over liquid oral preparations such as powders, capsules and tablets. If desired, capsules may be coated by standard aqueous or non-aqueous techniques. In addition to the dosage forms described above, the compounds of the invention may be administered by controlled release means and devices.

Pharmaceutical compositions of the present invention suitable for oral administration may be prepared as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient in powder or granular form or as a solution or suspension in an aqueous or nonaqueous liquid or in an oil-in-water or water-in-oil emulsion. Such compositions may be prepared by any of the methods known

in the art of pharmacy. In general, compositions are prepared by uniformly and intimately admixing the active ingredient with liquid carriers, finely divided solid carriers, or both and then, if necessary, shaping the product into the desired form. For example, a tablet may be prepared by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as a powder or granule optionally mixed with a binder, lubricant, inert diluent or surface active or dispersing agent. Molded tablets may be made by molding in a suitable machine, a mixture of the powdered compound moistened with an inert liquid diluent.

10

20

25

30

1.

It will be appreciated by one of ordinary skill in the art, however, that the most suitable route for administration will depend largely on the nature and severity of the condition being treated and on the nature of the active ingredient. As discussed above, the inventive therapeutics may be conveniently presented in unit dosage form and prepared by any of the methods well known in the art of pharmacy.

As discussed above, in one embodiment of the present invention, the inventive nalkenyl glycosides can be conjugated either directly or through a crosslinker to an appropriate carrier (e.g., KLH) to generate a synthetic tumor antigen. In general, a typical conjugation strategy that can be employed involves a reductive coupling of a glycoside which terminates in a glycoaldehyde, with the intended protein carrier, or lipid, presumably at the \varepsilon-amino acid residues of exposed lysines. (M.A. Bernstein; L.D. Hall, Carbohydr. Res. 1980, 78, C1; R.V. Lemieux Chem. Soc. Rev. 1978, 7, 423).

Thus, in another aspect, the present invention provides synthetic constructs, whereby novel antigenic structures, as described herein, are conjugated to carrier proteins, peptides or lipids. It will also be appreciated by one of ordinary skill in the art that, in the generation of a synthetic construct, more than one n-alkenyl moiety or glycopeptide moiety can ultimately be conjugated to a carrier protein to generate the synthetic vaccine. Thus, in addition to the conjugated glycopeptide structures as provided herein, constructs having the general structure as depicted below are also provided:

$$\begin{pmatrix}
A & C & CH_2 \\
A & CH_2
\end{pmatrix}_n & Carrier$$

wherein A is a carbohydrate domain having the structure:

10

15

5

wherein a, b, c, d, e, f, g, h, i, x, y and z are independently 0, 1, 2 or 3, with the proviso that x, y and z are not simultaneously 0; wherein R₀ is hydrogen, a linear or branched chain alkyl, acyl, arylalkyl or aryl group; wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈ and R₉ are each independently hydrogen, OH, ORⁱ, NH₂, NHCORⁱ, F, CH₂OH, CH₂ORⁱ, a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein Rⁱ is hydrogen, CHO, COORⁱⁱ, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group or a saccharide moiety having the structure:

5

10

wherein Y and Z are independently NH or O; wherein k, l, r, s, t, u, v and w are each independently 0, 1 or 2; wherein R₁₀, R₁₁, R₁₂, R₁₃, R₁₄ and R₁₅ are each independently hydrogen, OH, ORⁱⁱⁱ, NH₂, NHCORⁱⁱⁱ, F, CH₂OH, CH₂ORⁱⁱⁱ, or a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein R₁₆ is hydrogen, COOH, COORⁱⁱ, CONHRⁱⁱ, a substituted or unsubstituted linear or branched chain alkyl or aryl group; wherein Rⁱⁱⁱ is hydrogen, CHO, COOR^{iv}, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group; and wherein Rⁱⁱⁱ and R^{iv} are each independently H, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group;

wherein n is 0-8; wherein the carrier is a protein or lipid, including, but not limited to Bovine Serine Albumin, KLH and PamCys, wherein said protein or lipid is linked directly or though a crosslinker; and wherein m is in the range of 20-600. In certain preferred embodiments, n is 4. In certain other embodiments, m is in the range of 200-600. In still other preferred embodiments, the carbohydrate determinant is selected from the group consisting of Globo-H, KH-1, glycophorin, STN, (2,3)ST, N3, Tn, TF, 2,6-STn, and Le^y. In yet other preferred embodiments, the carbohydrate determinant is fucosyl GM1, which has the structure as depicted above, and as shown in Figure 1.

25

30

20

It will be appreciated that because certain of the inventive compounds produced terminate in an alkenyl linkage, in a typical protocol according to the present invention, conversion to an aldehyde is first required. Thus, in but one exemplary embodiment, an inventive synthetic globo-H tumor antigen is prepared from an n-alkenyl globo-H glycoside. As described in Example 2, this procedure involves exposing the n-alkenyl globo-H glycoside to oxidative conditions, in this case ozonolysis, followed by reductive

work-up to yield an aldehyde intermediate to generate a vaccine glycoconjugate. Subsequent hydrolytic carbohydrate analysis reveals approximately 350 carbohydrate residues/molecule of carrier protein, as described in Example 2.

In yet another example, a fucosyl GM1-KLH glycoconjugate is generated according to the method of the present invention, as discussed in Example 1. Notably, prior to conjugation studies, synthetic n-pentenyl fucosyl GM1 was shown to bind to monoclonal antibody F12 in ELISA and immune thin layer chromatography assays. Inhibition studies revealed that preincubation of F12 with antibody completely inhibits reactivity of natural fucosyl GM1, with the antibody. Clearly, the synthetic fucosyl GM1 pentenyl glycoside provides the antigenic epitope with which F12 reacts on SCLC cells.

10

15

20

25

30

35

Additionally, once a synthetic vaccine has been derivatized and characterized, mouse immunological studies can be performed to assess the potency and/or specificity of the novel tumor vaccines, as described in Example 4 herein.

Equivalents

The representative examples which follow are intended to help illustrate the invention, and are not intended to, nor should they be construed to, limit the scope of the invention. Indeed, various modifications of the invention and many further embodiments thereof, in addition to those shown and described herein, will become apparent to those skilled in the art from the full contents of this document, including the examples which follow and the references to the scientific and patent literature cited herein. In but one illustrative example, protecting groups play an important role in the synthesis of the carbohydrate domains and synthetic conjugates, as described herein; however it will be appreciated by one of ordinary skill in the art that the present invention encompasses the use of various alternate protecting groups known in the art. Those protecting groups used in the disclosure including the Examples below are merely illustrative.

It should further be appreciated that the contents of those cited references are incorporated herein by reference to help illustrate the state of the art. The following examples contain important additional information, exemplification and guidance which can be adapted to the practice of this invention in its various embodiments and the equivalents thereof.

5

10

15

20

25

30

(

Exemplification

A. Example 1: Synthesis of fucosyl GM1 pentenyl glycoside

1) Discussion of Synthesis:

As discussed above, in one aspect of the invention, the synthesis of fucosyl GM1 pentenyl glycoside is provided. In one embodiment of the present invention, this was achieved similarly to the methodology employed in the synthesis of MBr1 antigen, Globo-H (see, Park et al., J. Am. Chem. Soc. 1996, 118, 11488). For example, as shown in Figure 2, the synthesis of the ABC trisaccharide starting from the known protected lactal derivative 2, was first undertaken (Kwon, O.; Danishefsky. S.J. J. Am. Chem. Soc. 1998, 120, 1588). Selective sialylation of the C3' equatorial hydroxyl in 2 proceeded smoothly with phosphite donor 3 (Sim et al., J. Am. Chem. Soc. 1993, 115, 2260; Chappell et al., Tetrahedron 1997, 53, 11109) to yield the glycal 4 as the only observable isomer in 75% yield. In addition, propionitrile was employed as the solvent because of the necessity to perform the reaction at low temperatures. Use of elevated temperatures in acetonitrile as the solvent resulted in diminished anomeric selectivity, regioselectivity and lower chemical yields. The key DEF trisaccharide was synthesized as previously described in the Globo-H synthesis (Park et al. J. Am. Chem. Soc. 1996, 118, 11488). The requisite thioethyl donor 5 is shown in Figure 2. Based on previous experience, it was expected that this specific donor would favor β-glycosidation via sulfonamido participation under the close guidance of the "proximal hydroxyl" directing effect (see asterisk) (see also, Park et al. J. Am. Chem. Soc. 1996, 118, 11488; Kwon et al. J. Am. Chem. Soc. 1998, 120, 1588), and the results confirmed this expectation. In an experiment directed at "proof of principle", reaction of 5 with 5.0 equivalents of MeOTf (Lonn, H. Carbo. Res. 1985, 134, 105; Lonn, H. J. Carbohydr. Chem. 1987, 6, 301) in the presence of 4 gave the desired hexasaccharide 6 in 23 % yield, as shown in Figure 3. Although direct deprotection of this compound was not achieved to yield the desired compound, in an effort to find a hexasaccharide which was suitable for global deprotection, replacement of the reducing end glycal was considered.

replacement would also be potentially useful as a linker capable of being modified to allow for conjugation to a protein carrier or lipid.

10

15

20

25

30

35

In but one example, the use of a n-pentenyl glycoside was considered (For a review of n-pentenyl glycosides, see Fraser-Reid et al., Synlett, 1992, 927; Udodong et al. J. Am. Chem. Soc. 1993, 115, 7886; Merritt et al. J. Am. Chem. Soc. 1994, 116, 8334; Fraser-Reid et al. 1990, 55, 6068; Mootoo et al. J. Am. Chem. Soc. 1988, 110, 2662; Mootoo et al. J. Am. Chem. Soc. 1989, 111, 8540 and references therein). N-pentenyl glycosides are stable to a range of reaction conditions and reagents, but are readily activated for glycosidation reactions by treatment with a halogen oxidant. As a result of their stability and the neutral conditions required for their activation, pentenyl glycosides have been demonstrated to be valuable linkages for mechanistic and synthetic studies. Additionally, a terminal pentenyl group, or more generally a terminal alkenyl group, could also provide a handle for bioconjugation. Thus, in one embodiment, glycal 6a was subjected to epoxidation under standard procedures with 3,3-dimethyldioxirane (Figure 3). Reaction with pentenyl alcohol and anhydrous zinc chloride (Gordon et al. Carbohydrate Res. 1990, 206, 361) afforded the glycoside 7 in 65% yield. Indeed, with the pentenyl glycoside in place, global deprotection of 7 was possible. The sequence shown in Figure 3 furnished the peracetylated hexasaccharide lactone 8 in 46 % yield (5 steps). Removal of the acetates with sodium methoxide followed by saponification of the resulting methyl ester yielded the target, fucosyl GM1 pentenyl glycoside, 1b. The assignment of structure 1b was based on ¹H and ¹³C NMR analysis of 1b, in conjunction with characterization of intermediates en route to the final structure, and is supported by high resolution mass spectrometry.

In yet another embodiment, in an effort to produce significant quantities of this epitope for pre-clinical, and eventually clinical evaluation, a more efficient synthetic route was developed utilizing a glycoside at the reducing end at the acceptor, rather than a glycal. As shown in Figure 4, pentenyl lactoside was first investigated. For this purpose, lactose octaacetate was converted to the known bromide 9 (Reithal, Y. J. Am. Chem. Soc. 1952, 74, 4210; Dasgupta et al. Carbohydr. Res. 1994, 264, 155). Reaction of this compound with pentenyl alcohol under promotion by silver carbonate delivered the desired pentenyl glycoside, 10, on 100 g scale (Rodriguez, et al. Aust. J. Chem. 1990,

43, 665). An analogous coupling to produce 9 using silver triflate as promoter resulted in only a 17% yield of the desired product. Removal of the acetates yielded lactoside 11. Again, the C3' and C4' hydroxyl groups were engaged, this time as the dimethyl ketal 12. This reaction, as currently conducted, was accompanied by formation of minor amounts of 4,6-acetonide (Catelani et al. Carb. Res. 1988, 182, 297). Perbenzylation of 12 to give 13 followed by acetonide removal with aqueous acetic acid yielded the desired AB acceptor 14. Sialylation using phosphite donor 3 (Figure 2) proceeded in comparable yield to give trisaccharide acceptor, 15.

Finally, turning to the desired fucosyl GM1, coupling of donor 5 with a 2.0 molar excess of the acceptor 15 containing the pentenyl linker proceeded with MeOTf promotion (1.5 equivalents x 2) in excellent yield (70%; see Figure 5). Global deprotection under identical conditions as in Figure 4, yielded the characterized hexasaccharide 1b.

Attention was then directed to the final goal of developing a glycoconjugate. Synthetic 1b was subjected to conjugation to carrier protein KLH, as depicted in Figure 6. The protocol started with ozonolysis, thereby producing the uncharacterized aldehyde derivative. This step was followed by coupling to KLH using reductive amination under the agency of sodium cyanoborohydride. Presumably coupling of the carbohydrate had occurred with the \varepsilon-amino group of lysine residues in the KLH. Hydrolytic carbohydrate analysis revealed approximately 331 carbohydrate residues per molecule of KLH to generate 1c.

2) Experimentals

10

15

20

25

30

35

Peracetyl pentenyl-β-D-lactoside (10). To a cooled (ice bath) suspension of lactose octaacetate (100.0 g, 147.7 mmol), glacial acetic acid (30 mL) and acetic anhydride (30 mL) was added 100 mL of 30% HBr in AcOH dropwise over a period of 60 minutes. The reaction mixture stirred for 1 hour and the ice bath was removed. Upon stirring for an additional 2 hours at room temperature, the reaction became a homogeneous yellow solution. The solution was diluted with H₂O (1000 mL) and extracted with CHCl₃ (3x400 mL). The organic extracts were washed with H₂O (2x1000 mL), satd. NaHCO₃ (3x500 mL), dried over MgSO₄ and concentrated. The α-bromo

product 9 was azeotroped with anhydrous benzene and dried under high vacuum to yield

10

15

25

98.8 g (96%) of the lactosyl bromide which was used without further purification. To a suspension of Ag₂CO₃ (100 g, 362.6 mmol), freshly activated molecular sieves (15 g) and a crystal of I₂ in 400 mL CH₂Cl₂ was added pentenyl alcohol (5.0 equiv., 73.4 mL) and then the lactosyl bromide 9 (98.8 g, 141.4 mmol) in 400 mL of CH₂Cl₂. After stirring in the dark at room temperature for 16 hours, the reaction was filtered through a plug of celite with additional CH2Cl2 and concentrated to a yellow oil which was purified by flash column chromatography (10% EtOAc/hexanes->50% EtOAc/hexanes) to yield 74.7 g (75%) of the pentenyl lactoside 10 as a white foam. $[\alpha]^{22}$ D -48.9° (c 7.5, CHCl3); IR (film CHCl₃) 2941, 1751, 1369, 1224, 1054 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz) 85.60 (m, 1H), 5.17 (d, 1H, J=2.7 Hz), 5.02 (m, 1H), 4.93 (dd, 1H, J=7.9, 10.3 Hz), 4.85 (d, 1H, J=1.6 Hz), 4.78 (m, 2H), 4.71 (dd, 1H, J=9.6, 7.9 Hz), 4.30 (m, 3H), 3.93 (m, 3H), 3.66 (m, 3H), 3.45 (m, 1H), 3.30 (m, 1H), 1.98 (s, 3H), 1.94 (s, 3H), 1.91 (m, 2H), 1.89 (s, 3H), 1.88 (s, 6H, 2xCH₃), 1.87 (s, 3H), 1.79 (s, 3H), 1.49 (m, 2H); ¹³C NMR (CDCl₃, 100 MHz) 8 170.33, 170.28, 170.09, 170.00, 169.74, 69.54, 169.01, 137.72, 115.00, 101.01, 100.51, 76.27, 72.76, 72.48, 71.64, 70.94, 70.58, 69.23, 69.01, 66.52, 61.97, 60.73, 29.75, 28.49, 20.80, 20.75, 20.64, 20.57, 20.45; HRMS (FAB) cald. for C₃₁H₄₄O₁₈Na [M+Na]⁺ 727.2425, found 727.2418. Pent-4-enyl 3', 4'-O-acetonide-β-D-lactoside (12). Peracteylated pentenyl lactoside, 10,

(18.2 g, 25.8 mmol) was dissolved in 300 mL of anhydrous MeOH and 2.0 mL of NaOMe (25% in MeOH) was added. The reaction stirred at rt for 16 hours and was neutralized with Dowex-H⁺ (pH 5-6). The reaction was filtered with additional MeOH and concentrated to a white solid (10.6 g, quantitative) which was used without further purification. Pentenyl- β -D-lactoside 11: 1 H NMR (D₂O, 400 MHz) δ 5.81 (m, 1H), 5.00 (dd, 1H, J=17.3, 1.9 Hz), 4.92 (dd, 1H, J=8.9 Hz), 4.74 (m, 1H), 4.39 (d, 1H, J=8.0 Hz), 4.35 (d, 1H, J=7.8 Hz), 3.72-3.42 (m, 12H), 3.21 (m, 1H0, 2.06 (m, 2H), 1.63 (m, 2H); 13 C NMR (D₂O, 100 MHz) δ 141.27, 117.31, 105.42, 104.54, 80.85, 77.84, 77.24, 76.92, 75.33, 75.00, 73.44, 72.47, 71.03, 63.52, 62.56, 31.83, 30.48.

To pentenyl lactoside 11 (10.6 g, 25.8 mmol) was added 200 mL acetone, 26 mL of dimethoxypropane and p-toluenesulfonic acid (491 mg, 0.1 equiv.). The suspension stirred at room temperature overnight at which point the reaction was homogeneous. The reaction was concentrated and purified by flash column chromatography (100% EtOAc -> 2%MeOH in EtOAc) to give 5.1 g (44%) of the 3,4-acetonide as a white solid and 1.27 g of the 4,6-acetonide as a white solid. 3,4-acetonide, 12: [\alpha]^{22} \dagger 79.0° (0.96c. MeOH): IR 3422, 2980, 2933, 2870, 1242, 1073 cm⁻¹; ¹H NMR (MeOH, 400 MHz) δ 5.83 (m, 1H), 5.00 (dd, 1H, J=17.1, 3.4 Hz), 4.92 (dd, 1H, J=10.2, 2.0 Hz), 4.34 (d, 1H, J=8.2 Hz), 4.25 (d, 1H, J=7.8 Hz), 4.17 (dd, 1H, J=5.5, 2.1 Hz), 4.02 (dd, 1H, J=7.2, 5.5 Hz), 3.91 (m, 3H), 3.88-3.73 (m, 5H), 3.55-3.47 (m, 3H), 3.42 (m, 1H), 3.38 (m, 1H), 3.21 (m, 1H), 2.13 (m, 2H), 1.67 (m, 2H), 1.45 (s, 3H), 1.30 (m, 3H); ¹³C NMR (MeOH, 100 MHz) δ 139.42, 115.20, 111.04 (O-C-O), 104.16, 104.09, 80.94, 80.77, 76.29, 76.25, 75.27, 75.00, 74.76, 74.39, 62.36, 61.82, 31.18 (CH₃), 30.02, 28.41, 26.51 (CH₃); HRMS (Fab) cald. for $C_{20}H_{34}O_{11}Na [M+Na]^{+}$ 473.1998, found 473.1985. 4,6-acetonide: $[\alpha]^{22}D$ -216.0 ° (c 1.14, MeOH); IR 3364, 2926, 2870, 1380 cm⁻¹; ¹H NMR (MeOH, 400 MHz) δ 5.79 (m, 1H), 4.98 (dd, 1H, J=17.0, 1.8), 4.90 (dd, 1H, J=10.2, 1.0), 4.35 (d, 1H, J-7.8 Hz), 4.24 (d, 1H, J=7.8 Hz), 4.13 (m, 2H), 3.86-3.82 (m, 3H), 3.76 (dd, 1H, J=12.9, 1.4 Hz), 3.61-3.49 (m, 5H), 3.44 (s, 1H), 3.35 (m, 1H), 3.19 (t, 1H), 2.11 (m, 2H), 1.66 (m, 2H), 1.43 (s, 3H), 1.35 (s, 3H); ¹³C NMR (MeOH, 100 MHz) δ 139.48, 115.20, 104.68. 104.26, 100.17 (O-C-O), 79.86, 76.40, 76.35, 74.81, 73.34, 71.46, 70.25, 69.95, 68.04. 63.67, 61.69, 31.23 (CH₃), 30.08, 29.56, 18.69 (CH₃).

10

25

Pent-4-enyl 2, 3, 6, 2', 6'-penta-O-benzyl-3', 4'-O-acetonide-β-D-lactoside (13). The acetonide 12 (3.40 g, 7.54 mmol) was azeotroped with anhydrous benzene, dissolved in anhydrous DMF (60 mL, 0.12 M) and cooled to 0 °C. Benzyl bromide (passed through basic alumina, 10.0 equiv. 8.97 mL), was added followed by solid NaH (95%, 7.5 equiv., 1.76 g) in one portion. The reaction was allowed to warm to room temperature overnight and then poured into ice cold H₂O (500 mL) and extracted with CHCl₃ (200 mL, 2x100mL). The organic extracts were washed with brine (500 mL), dried over MgSO₄ and concentrated to a yellow oil which was purified by flash column chromatography

(5% EtOAc/hexanes -> 20% EtOAc/hexanes) to yield 5.70 g (84%) of the product as a viscous oil. [α]²²D 196.0° (1.09c, CHCl3); IR (film CHCl3) 3062, 3029, 2868, 1367, 1093, 1055 cm⁻¹; ¹H NMR (CDCl3, 400 MHz) δ 7.39-7.20 (m, 25H), 5.78 (m, 1H), 4.98 (dd, 1H, J=17.1, 3.4 Hz), 4.93 (dd, 1H, J=10.2, 2.0 Hz), 4.89 (d, 1H, J=10.6 Hz), 4.86 (d, 1H, J=10.9 Hz), 4.75 (d, 1H, J=11.7 Hz), 4.70 (d, 1H, J=10.6 Hz), 4.68 (d, 1H, J=10.8 Hz), 4.63 (d, 1H, 11.8 Hz), 4.53 (d, 1H, J=12.1 Hz), 4.46 (d, 1H, J=12.1 Hz), 4.39 (d, 1H, J=6.3 Hz), 4.36 (d, 1H, J=2.0 Hz), 4.34 (d, 1H, J=7.8 Hz), 4.28 (d, 1H, J=12.0 Hz), 4.07 (dd, 1H, J=5.5, 1.4 Hz), 3.99 (m, 1H), 3.91 (m, 2H), 3.76 (dd, 1H, J=11.9, 4.2 Hz), 3.70 (dd, 1H, J=10.8, 1.6 Hz), 3.65 (m, 2H), 3.55-3.47 (m, 3H), 3.35 (m, 2H), 3.30 (dd, 1H, J=7.9, 0.8 Hz), 2.31 (m, 2H), 1.72 (m, 2H), 1.36 (s, 3H), 1.31 (s, 3H); ¹³C NMR (CDCl3, 100 MHz) δ 138.92, 138.54, 138.43, 138.31, 138.19, 138.00, 128.22, 128.16, 128.08, 127.99, 127.95, 127.77, 127.63, 127.49, 127.40, 127.34, 127.20, 114.83, 109.66, 103.56, 101.76, 82.89, 81.75, 80.53, 79.26, 76.30, 75.32, 74.96, 74.91, 73.51, 73.28, 73.12, 73.08, 71.86, 69.16, 68.82, 68.18, 30.15, 28.87, 27.89, 26.34; HRMS (FAB) cald. for [M+Na]⁺ C₅₅H₆₄O₁₁Na 923.4346, found 923.4330.

Pent-4-enyl 2, 3, 6, 2', 6'-penta-*O*-benzyl-β-D-lactoside (14). The acetonide 13 (5.7 g, 6.32 mmol) was dissolved in 80% AcOH in H₂O (60 mL) and heated to 75 °C for 3 hours. The reaction was cooled to rt, diluted with H₂O (500 mL) and extracted with CHCl₃ (200 mL, 2x100 mL). The organic extracts were washed with H₂O (500 mL), satd. NaHCO₃ (3x300 mL), dried over MgSO₄ and concentrated to an oil which was purified by flash column chromatography (25% EtOAc/hexanes) to yield 5.21 g (96%) of a white solid. [α]²²D 194.1° (1.13c, CHCl₃); IR (film CHCl₃) 3444, 3028, 2868, 1091, 1058 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz) δ 7.41-7.20 (m, 25H), 5.74 (m, 1H), 4.96-4.88 (m, 3H), 4.82 (d, 1H, J=10.9 Hz), 4.72 (d, 1H, J=11.4 Hz), 4.70 (d, 1H, J=10.9 Hz), 4.64 (d, 1H, J=10.9 Hz), 4.58 (d, 1H, J=11.6 Hz), 4.52 (d, 1H, J=10.9), 4.38-4.28 (m, 5H), 3.93-3.85 (m, 3H), 3.71 (m, 2H), 3.55-3.40 (m, 4H), 3.36 (m, 3H), 3.28 (m, 2H), 2.48 (d, OH, 1H, J=3.2 Hz), 2.40 (s, OH, 1H), 2.09 (m, 2H), 1.66 (m, 2H); ¹³C NMR (CDCl₃, 100 MHz) δ 139.09, 138.53, 139.29, 138.17, 138.01, 137.91, 128.44, 128.35, 128.24, 127.89, 127.82, 127.77, 127.63, 127.55, 127.19, 114.86, 103.57, 102.53, 82.76, 81.71,

79.96, 76.57, 75.17, 75.02, 74.89, 74.80, 73.45, 73.40, 73.12, 72.79, 69.22, 68.71, 68.60, 68.24, 30.17, 28.88; HRMS (FAB) cald. for C₅₂H₆₀O₁₁Na [M+Na]⁺883.4033, found 883.4017.

(

10

20

25

30

Trisaccharide 15. The phosphite donor 3 (1.0 g, 1.35 mmmol) and lactosyl acceptor 14 (2.5 g. 2.90 mmol) were combined, azeotroped with anhydrous benzene and placed under high vacuum for 2 hours. The mixture was dissolved in anhydrous CH3CH2CN (distilled from CaH₂), freshly activated molecular sieves were added and the reaction cooled to -40 °C. A portion of TMSOTf (0.1 equiv., 27µL) was added and the reaction was allowed to stir at -40 °C overnight. The reaction was then warmed to -30 °C and another 0.1 equivalent of TMSOTf was added. Upon stirring for an additional 2 hours at -30 °C, the reaction was quenched by the addition of solid NaHCO3 and was filtered through a plug of celite with the aid of EtOAc. The organic layer was washed with satd. NaHCO₃ (2x400 mL) and dried over MgSO₄. Evaporation of organic layer gave a cloudy oil which was subjected to flash column chromatography using careful gradient elution in order to recover acceptor and product trisaccharide (20% EtOAc/hexanes -> 75% EtOAc/hexanes). The product (1.35g, 75%) was obtained as a white foam and 0.95 g of starting acceptor was recovered. : $[\alpha]^{22}D$ 2.38° (c 1.30, CHCl₃); IR (film CHCl₃) 3106, 2866, 1744, 1689, 1368, 1222, 1055 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz) δ 7.40-7.17 (m, 25H), 5.79 (m, 1H), 5.38 (m, 1H), 5.27 (dd, 1H, J=8.0, 2.0 Hz), 5.08 (d, 1H, J=10.0 Hz), 4.95 (m, 3H), 4.86 (d, 1H, J=10.9 Hz), 4.75 (d, 1H, J=5.7 Hz), 4.72 (d, 1H, J=10.8 Hz), 4.68 (d, 1H, J=11.0 Hz), 4.56 (d, 1H, J=11.9 Hz), 4.54 (d, 1H, J=7.6 Hz), 4.44 (d, 1H, J=12.2 Hz), 4.39 (m, 1H), 4.32-4.25 (m, 3H), 4.06-3.88 (m, 6H), 3.79 (m, 2H), 3.72 (s, 3H), 3.65 (m, 3H), 3.54-3.44 (m, 5H), 3.35 (m, 2H), 2.66 (d, OH, 1H, J=3.3 Hz), 2.47 (dd, 1H, J=13.0, 4.7 Hz), 2.12 (m, 2H), 2.06 (s, 3H), 2.02 (m, 1H), 1.98 (s, 3H), 1.95 (s, 3H),1.85 (s, 3H), 1.83 (s, 3H), 1.71 (s, 3H); 13 C NMR (CDCl₃, 100 MHz) δ 170.77, 170.53, 170.23, 169.92, 169.87, 168.32, 139.09, 138.90, 138.61, 138.45, 138.34, 138.05, 128.27, 128.21, 137.99, 127.51, 127.42, 127.11, 114.81, 103.48, 102.29, 98.32, 82.90, 81.80, 78.37, 76.50, 76.30, 75.31, 75.01, 74.89, 74.82, 73.23, 72.97, 72.66, 72.37, 69.16, 69.03, 68.69, 68.43, 68.36, 67.81, 67.08, 62.21, 52.99, 49.17, 36.41, 30.17, 28.89, 23.11, 21.08, 20.77, 60.67, 60.47; HRMS (FAB) cald. for $C_{72}H_{87}NO_{23}Na$ (M+Na⁺) 1356.5566, found 1356.5557.

10

15

25

Hexasaccharide 7 (R=Bn). The thioethyl donor 5 (311 mg, 0.243 mmol) and acceptor 15 (627 mg, 0.487 mmol) were combined, azeotroped with anhydrous benzene (5x5 mL) and placed under high vacuum for 5 hours. The mixture was then dissolved in 1.6 mL CH2Cl2 and 3.2 mL Et2O (0.05M total), treated with freshly prepared molecular sieves and cooled to 0 °C. Methyl triflate (1.5 equiv., 41 µL) was added in one portion and the reaction stirred at 0 °C overnight. In the morning, another 20 µL of MeOTf was added and the reaction was allowed to stir for an additional 2 hours at 5 °C. The reaction was quenched by the addition of solid NaHCO3, filtered through celite with EtOAc, concentrated and purified by flash column chromatography (gradient elution 25% EtOAc/hexanes -> 50% -> 75% EtOAc/hexanes) to give 437 mg (70%) of hexasaccharide as a white foam and 300 mg of recovered trisaccharide acceptor: [\alpha]^{22}D -29.4° (c 3.25, CHCl3); IR (film CHCl3) 3285, 3028, 2940, 2865, 1794, 1749, 1690, 1220, 1090 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz) δ 7.74 (d, 2H, J=7.5 Hz), 7.34-7.08 (m, 43H), 5.75 (m, 1H), 5.52 (d, 1H, J=4.7 Hz), 5.29 (app s, 1H), 5.23 (dd, 1H, J=9.5, 1.4 H2), 5.15 (m, 1H), 5.02 (d, 1H, J=9.8 H2)4.97-4.87 (m, 5H), 4.84 (d, 1H, J=10.9 Hz), 4.81-4.70 (m, 5H), 4.63 (d, 1H, J=11.6 Hz), 4.57 (m, 3H), 4.44 (d, 1H, J=7.2 Hz), 4.40 (d, 1H, J=12.2 Hz), 4.30 (d, 1H, J=7.8 Hz), 4.10 (m, 2H), 3.98-3.81 (m, 12H), 3.82 (s, 3H), 3.78-3.68 (m, 7H), 3.64-3.45 (m, 8H), 3.27 (m, 3H), 3.17 (dd, 1H), 2.80 (d, OH, 1H, J=2.1 Hz), 2.19 (dd, 1H, J=13.0, 4.5 Hz), 2.10 (m, 3H), 2.01 (s, 3H), 1.92 (s, 3H), 1.88 (s, 3H), 1.82 (s, 3H), 1.81 (s, 3H), 1.68 (m, 2H), 1.08 (d, 3H, J=5.4 Hz), 1.00-0.92 (m, 42H); ¹³C NMR (CDCl₃, 100 MHz) δ 170.61, 170.34, 170.26, 169.66, 167.78, 155.48, 138.95, 138.65, 138.63, 138.56, 138.42, 138.38, 138.27, 138.05, 132.17, 129.02, 128.59, 128.46, 128.18, 128.05, 127.91, 127.63, 127.51, 127.24, 127.09, 114.80, 103.42, 102.76, 102.45, 100.16, 99.58, 98.76, 82.87, 81.53, 79.06, 77.32, 77.24, 77.16, 75.12, 75.07, 74.95, 74.80, 73.92, 73.27, 73.04, 72.93, 72.19, 69.23, 69.14, 69.09, 67.89, 67.53, 61.76, 61.58, 61.12, 56.39, 53.60, 49.19, 35.36, 30.17, 28.89, 23.13, 20.97, 20.75, 20.62, 20.53,

17.85, 17.53, 17.33, 16.72, 11.80, 11.74; HRMS (FAB) cald. for C₁₃₆H₁₇₈N₂O₃₉SSi₂ (M+Na⁺) 2574.1163, found 2574.1130.

10

15

20

25

30

Compound 1b. To a solution of the hexasaccharide (130 mg, 0.0509 mmol) in THF (2.0 mL) was added glacial AcOH (10.0 equiv., 29 μL) and TBAF (1.0 M THF, 10.0 equiv., 0.509 mL). The reaction stirred at rt overnight, was poured into ice water and extracted with EtOAc (3x50 mL). The organic extracts were washed with satd. NaHCO3 (50 mL) and brine (50 mL), dried over MgSO4 and concentrated to an oil which was purified through a short plug of silica gel with EtOAc. The resulting triol was dissolved in anhydrous MeOH (2.5 mL) and sodium methoxide was added (0.250 mL of a 25% solution in MeOH). The reaction stirred at rt for 18 hours and then 0.5 mL of THF and 0.5 mL of H2O were added. Stirring at rt for an additional 24 hours was followed by neutralization with Dowex-H+, filtration with MeOH washings and concentration. The crude material was allowed to dry under high vacuum for 1 day. To the resulting white solid was added THF (0.5 mL) and condensed liquid NH3 (~10 mL) at -78 °C. Sodium (~50 mg) was added and the resulting blue solution stirred at -78 °C for 1.5 hours. The reaction was quenched with anhydrous MeOH (~5 mL), brought to rt and concentrated with a stream of dry N2 to a volume of ~ 2 mL. The reaction was neutralized with Dowex-H+, filtered with MeOH washings and concnetrated to a white solid. The white solid was dissolved in 1.0 mL pyridine and 1.0 mL CH2Cl2 and cooled to 0 °C. A crystal of DMAP was added followed by acetic anhydride (1.0 mL). The ice bath was removed and the reaction stirred at rt overnight. Concentration followed by purification by flash column chromatography (gradient elution 75% EtOAc/hexanes -> 100% EtOAc -:> 5% MeOH/EtOAc) gave 44 mg (46%) of 8 as a white solid: ¹H NMR (MeOH, 400 MHz) δ 8.02 (d, 1H, J=9.9 Hz), 7.87 (d, 1H, J=9.2 Hz), 5.76 (m, 1H), 5.49 (m, 1H), 5.39 (d, 1H, J=2.9 Hz), 5.34-5.31 (m, 2H), 5.22 (d, 1H, J=3.4 Hz), 5.19 (d, 1H, J=4.1 Hz), 5.17 (d, 1H, J=3.5 Hz), 5.12-5.05 (m, 3H), 4.97 (dd, 1H, J=16.8, 1.7 Hz), 4.91 (dd, 1H, J=10.0, 1.7 Hz), 4.81-4.75 (m, 3H), 4.65-4.60 (m, 2H), 4.52 (d, 1H, J=7.9 Hz), 4.48-4.44 (m, 2H), 4.37 (dd, 1H, J=10.0, 2.5 Hz), 4.28 (dd, 1H, J=12.5, 2.4 Hz), 4.22-4.18 (m, 2H), 4.14-3.99 (m, 9H), 3.96-3.92 (m, 2H), 3.89 (d, 1H, J=2.9 Hz), 3.88-3.77 (m, 4H), 3.72-3.62 (m, 3H), 3.51-3.45 (m, 1H), 2.74 (dd, 1H, J=11.3, 4.5 Hz), 2.19 (s, 3H), 2.13 (s, 3H), 2.11 (s, 3H), 2.10 (s, 3H), 2.09 (s, 3H), 2.08 (s, 3H), 2.06 (s, 3H), 2.05 (s, 3H), 2.02 (s, 3H), 2.01 (s, 3H), 2.00 (s, 3H), 1.99 (s, 3H), 1.98 (s, 3H), 1.97 (s, 3H), 1.95 (s, 3H), 1.94 (s, 3H), .91 (s, 3H), .180 (s, 3H), 1.61 (m, 2H), 1.14 (d, 3H, J=6.4 Hz), 3 protons burried beneath acetates (2 Pn, 1 C3ax); ¹³C NMR (MeOH, 100 MHz) δ 174.64, 173.64, 172.98, 172.89, 172.63, 172.56, 172.48, 172.44, 172.34, 172.27, 172.04, 171.99, 171.76, 171.73, 171.62, 171.35, 171.25, 139.23, 115.47, 104.62, 103.26, 101.86, 101.63, 100.78, 97.31, 78.22, 76.53, 75.08, 74.69, 74.29, 73.91, 73.53, 72.94, 72.71, 72.56, 72.16, 72.06, 71.89, 71.74, 70.19, 69.87, 69.33, 69.11, 68.92, 65.96, 65.65, 63.68, 63.52, 62.69, 54.01, 53.09, 50.60, 40.19, 31.09, 29.96, 24.17, 24.06, 22.73, 21.76, 21.59, 21.46, 21.20, 21.06, 20.89, 20.75, 20.63, 20.55, 16.52.

The peracetate (40 mg) was dissolved in anhydrous MeOH (2.0 mL) and 150 µL of 15 sodium methoxide was added (25% solution in MeOH). The reaction stirred at rt for 18 hours and then 0.5 mL of THF and 0.5 mL of H2O was added. The reaction stirred for another 24 hours at rt. Neutralization with Dowex-H+ (~pH 6-7) was followed by filtration with MeOH washings, concentration and purification using P-2 Gel (H2O elutent) to yield 24 mg (96%) of a white solid: IR 3346, 2940, 2882, 1657, 1620, 1376, 20 1069 cm⁻¹; ¹H NMR (D₂O, 400 MHz) δ 5.86 (m, 1H), 5.18 (d, 1H, J=4.0 Hz), 5.04 (dd, 1H, J=17.22, 1.7 Hz), 4.97 (dd, 1H, J=10.6 Hz), 4.63 (d, 1H, J=7.6 Hz), 4.57 (d, 1H, J=7.7), 4.46 (d, 1H, J=7.9 Hz), 4.43 (d, 1H, J=8.1 Hz), 4.15 (m, 1H), 4.09-4.02 (m, 3H), 3.94-3.84 (m, 5H), 3.80-3.63 (m, 18H), 3.60-3.53 (m, 6H), 3.47 (dd, 1H, J=10.3, 1.8), 3.32 (t, 1H), 3.26 (t, 2H), 2.62 (dd, 1H, J=13.4, 4.3 Hz), 2.09 (m, 2H), 1.98 (s, 6H), 1.86 25 (m, 1H), 1.67 (m, 2H), 1.15 (d, 3H, J=6.5 Hz); 13 C NMR (D₂O, 100 MHz) δ 176.29, 175.43, 175.16, 139.97, 115.99, 104.38, 103.77, 103.30, 103.22, 102.25, 100.35, 79.67, 78.12, 77.65, 77.03, 76.06, 75.94, 75.62, 75.44, 75.24, 74.85, 74.19, 74.01, 73.45, 73.01, 71.15, 70.72, 70.32, 69.87, 69.64, 69.25, 67.93, 64.01, 62.29, 62.07, 61.63, 61.29, 52.79, 52.70, 50.04, 38.45, 30.53, 29.17, 23.89, 23.23, 16.53; HRMS (FAB) cald. for 30 $C_{48}H_{79}N_2O_{33}Na_2 [M-H+2Na]^+ 1257.4360$, found 1257.4337.

Glycal hexasaccharide 6a. The thioethyl donor 5 (120 mg, 0.0938 mmol) and acceptor 4 (122 mg, 0.108 mmol) were combined, azeotroped with anhydrous benzene (5x5mL) and placed under high vacuum overnight. The mixture was dissloved in a 2:1 mixture of

15

25

()

Et2O:CH2Cl2 (2.7 mL total), molecular sieves were added and the mixture stirred at rt for 1 h. The reaction was cooled to 0 °C and 1.0 equiv. of MeOTf (0.020 mL) was added. After 4 hours at 0 °C another equivalent of MeOTf was added (0.020mL) and the reaction continued to stir for another 4 h at 10 °C. The reaction was quenched with solid NaHCO3, filtered through celite with additional EtOAc (100ml) and concentrated. The resulting mixture was purified by flash column chromatography to give 50 mg (23%) of the hexasaccharide glycal 6 and 85 mg of starting acceptor, 4: R_f 0.35 (66% EtOAc/hexanes); ¹H NMR (500 MHz, C₆D₆) δ 8.31 (d, 2H), 7.62 (d, 2H), 7.52 (m, 4H), 7.45 (d, 2H), 7.40-7.15 (m, 31H), 6.47 (d, 1H, J=6.3 Hz), 6.28 (apparent s, 1H), 6.09 (d, 1H, J=3.8 Hz), 5.72 (m, 1H), 5.55 (dd, 1H, J=9.3, 1.2 Hz), 5.51 (d, 1H, J=3.5 Hz), 5.22 (d, 1H, J=10.8 Hz), 5.15 (s, 1H), 5.13-5.06 (m, 3H), 5.05 (d, 1H, J=8.1 Hz), 5.02 (m, 1H), 4.98 (d, 1H, J=10.8 Hz), 4.85 (d, 1H, J=10.6 Hz), 4.82 (d, 1H, J=9.4 Hz), 4.73-4.66 (m, 8H), 4.55-4.34 (m, 10H), 4.38-4.32 (m, 5H), 4.30 (d, 1H), 4.18 (s, 3H), 4.21-4.12 (m, 6H), 4.06 (m, 2H), 3.99 (m, 4H), 3.85 (d, 1H), 3.74 (dd, 1H), 3.61 (m, 2H), 3.52 (t, 1H), 2.63 (dd, 1H, J=13.9, 5.0), 2.48 (dd, 1H, J=13.4 Hz), 2.35 (s, 3H), 2.01 (s, 3H), 1.98 (s, 3H), 1.72 (s, 3H), 1.64 (s, 3H), 1.57 (d, 3H, J=6.3), 1.31-1.20 (m, 42H); ¹³C NMR (100 MHz, CDCl₃) δ 169.71, 169.39, 169.18, 168.70, 168.12, 166.99, 154.75, 143.47, 137.81, 137.71, 137.51, 137.42, 137.07, 131.65, 128.25, 127.52, 128.32, 127.26, 127.23, 127.19, 127.10, 126.98, 126.91, 126.83, 126.73, 126.62, 126.53, 126.36, 126.29, 101.67, 101.35, 98.69, 98.32, 98.26, 97.33, 80.48, 78.05, 77.06, 76.20, 75.50, 74.64, 74.22, 73.87, 73.49, 72.90, 72.38, 72.26, 71.93, 71.47, 71.20, 70.34, 70.17, 69.99, 69.13, 68.62, 68.10, 67.92, 67.01, 66.88, 66.68, 65.52, 60.92, 60.61, 55.51, 52.59, 48.31, 34.87, 28.68, 22.19, 19.95, 19.77, 19.68, 19.59, 16.93, 16.88, 15.79, 10.86, 10.78; HRMS (FAB) cald. for $C_{124}H_{162}N_2O_{37}Si_2SNa [M+Na]^{\dagger} 2382.0013$, found 2382.0001.

Imido-hexasaccharide 6b Performing the above reaction with 10 equiv. MeOTf added in one portion, under otherwise identical conditions yields 28% of the following compound, which is much less polar than the parent N-acetylated hexasaccharide 6a. R_f 0.35 (25% EtOAc/hexanes); ¹H NMR (500 MHz, C₆D₆) δ 8.31 (d, 2H), 7.66 (d, 2H), 7.53 (t, 4H), 7.48 (d, 2H), 7.42-7.16 (m, 31H), 6.46 (d, 1H), 6.21 (app s, 1H), 6.15 (d, 1H, J=4.3 Hz), 5.81 (d, 1H, J=9.2 Hz), 5.72 (dt, 1H, J=12.8, 2.4 Hz), 5.40 (m, 1H), 5.38 (d,

1H, J=3.5 Hz), 5.20 (d, 1H, J=10.2 Hz), 5.12 (t, 2H), 5.00 (m, 3H), 4.84 (d, 1H, J-6.2 Hz), 4.81 (d, 1H, J=4.5 Hz), 4.73 (m, 2H), 4.70 (m, 2H), 4.67 (d, 1H, J=2.6 Hz), 4.65 (m, 1H), 4.59 (m, 3H), 4.53-4.46 (m, 6H), 4.40 (m, 5H), 4.36 (d, 1H, J=3.1 Hz), 4.30 (d, 1H, J=3.4 Hz), 4.26 (m, 3H), 4.23 (app s, 1H), 4.20 (m, 3H), 4.11 (m, 2H), 4.04 (d, 1H, J=5.9 Hz), 3.99 (s, 3H), 3.92 (d, 1H, J=3.2 Hz), 3.87 (d, 1H, J=2.9 Hz), 3.82 (d, 1H, J=6.5 Hz), 3.70 (m, 1H), 3.64 (s, 3H), 3.60 (d, 1H), 3.28 (t, 1H), 2.94 (dd, 1H, J=13.7, 4.5 Hz), 2.36 (t, 1H, J=13.3 Hz), 2.14 (s, 3H), 1.91 (s, 3H), 1.83 (s, 3H), 1.81 (s, 3H), 1.60 (s, 3H), 1.53 (d, 3H, J=6.5 Hz), 1.32-1.23 (m, 42H); 13 C NMR (100 MHz, CHCl₃) δ 170.43, 169.30, 169.20, 168.98, 168.03, 164.74, 155.82, 144.74, 139.09, 138.75, 138.52, 138.48, 138.40, 138.39, 138.25, 138.17, 132.56, 129.22, 128.85, 128.39, 128.35, 128.30, 128.25, 128.01, 127.79, 127.71, 127.60, 127.55, 127.50, 127.48, 127.34, 102.57, 102.24, 99.69, 99.11, 98.25, 81.35, 79.09, 87.22, 75.64, 75.40, 74.90, 74.60, 74.15, 73.95, 73.50, 73.33, 72.94, 72.84, 72.52, 71.37, 71.17, 70.47, 70.17, 69.66, 69.05, 68.47, 68.11, 67.96, 67.71, 67.55, 61.91, 61.54, 61.05, 57.70, 56.50, 53.65, 52.75, 31.94, 29.71, 21.70, 20.97, 20.89, 20.64, 20.46, 20.44, 17.57, 16.81, 15.38, 14.13, 11.89, 11.80; LRMS (FAB) $C_{125}H_{164}N_2O_{37}SSi_2Na 2373 [M+Na]^{+}$.

3) Conjugation studies:

20

25

35

As described herein and as shown in Figure 6, the pentenyl group in FucGM1 was converted to an aldehyde group by ozonolysis and linked to -NH2 groups of KLH by reductive amination method in the presence of sodium cyanoborohydride as described for globo H (see, Ragupathi G, Park TK, Zhang S, Kim IJ, Graeber L, Adluri S, Lloyd KO, Danishefsky SJ and Livingston PO. Immunization of mice with conjugates of fully synthetic hexasaccharide globo H results in antibody against human cancer cells: a combined chemical immunological approach to the fashioning of an anticancer vaccine. Angewandte Chem. Int. Ed Engl. 36: 125-128. 1997.). In the case of cross-linker method the aldehyde group obtained through ozonolysis was first reacted with hydrazide of MMCCH (4-(maleimidomethyl) cyclohexane-1-carboxyl hydrazide) and reacted with thiolated KLH as described in Ragupathi G, Koganty RR, Qiu D, Lloyd KO and Livingston PO. A novel and efficient method for synthetic carbohydrate conjugate vaccine preparation: Synthesis of sialyl Tn-KLH conjugate using a 4-(4-N-

maleimidomethyl) cyclohexane-1-carboxyl hydrazide (MMCCH) linker arm. Glycoconjugate J., 15: 217-221, 1998). For example, 4 mg of FucGM1 pentenyl glycoside in methanol was stirred at -78°C in a dry-ice/ethanol bath and ozone gas was passed through the solution for 10 min under vigorous stirring. The excess of ozone was then displaced with nitrogen over a period of 5 min. Methyl sulfide (100 µl) was added and the reaction mixture stirred at room temperature for 2 hours and distributed equally in two vials. The solvent was removed under a stream of nitrogen. The resulting white solid was used directly in the subsequent conjugation steps.

a) Direct Conjugation of FucGM1-aldehyde with KLH:

(

10

15

20

25

30

Two mg FucGM1-aldehyde was dissolved in 1 ml of 0.1M phosphate buffered saline (PBS) pH 7.2 and 4 mg of KLH in PBS. Two mg sodium cyanoborohydride was added and the mixture incubated under gentle agitation at 37°C for 48 h. After 16 h, an additional 1.0 mg sodium cyanoborohydride was added and the incubation continued. The unreacted FucGM1 aldehyde was removed completely with multiple washes using a Amicon Centriprep with molecular weight cut-off value 30000 dalton, with 6-7 changes of PBS at 4°C.

b) Conjugation of FucGM1-aldehyde through MMCCH to thiolated KLH: Preparation of FucGM1-MMCCH

Two mg of FucGM1-aldehyde was dissolved in 1 ml of 0.1M sodium acetate buffer pH 5.5, and 4 mg of MMCCH in 100 µl of dimethyl sulfoxide (DMSO) was added. The reaction mixture was incubated at room temperature for 15 min with gentle stirring. At the end of 15 min 2 mg of solid sodium cyanoborohydride was added and the incubation continued at room temperature for 2 h. Unreacted MMCCH was removed in a Sephadex G10 column equilibrated previously with 0.1 M sodium phosphate buffer pH 6.0 containing 5 mM EDTA and eluted with the same buffer. The fractions positive for FucGM1 by TLC with orcinol were combined.

Addition of sulfhydryl groups to KLH

15

20

25

30

35

2-Iminothiolane (2 mg) dissolved in thiolation buffer (50 mM triethanolamine, 0.15 M NaCl, 5 mM EDTA, pH 8.0) was added to 4 mg of KLH and incubated with stirring at room temperature for 2 h. Unreacted 2-iminothiolane was removed by Sephadex G15 column equilibrated previously with 0.1 M sodium phosphate buffer pH 7.2 containing 5 mM EDTA and eluted with the same buffer. Fractions positive for KLH with BioRad protein assay dye reagent were combined. A small portion was used to estimate sulfhydryl groups in the thiolated KLH using Ellman's reagents and cysteine as standard as described earlier (Riddles PW, Blackeley RL, Zemer B Ellman's reagent: 5,5'-dithiobis(2-nitrobenzoic acid)--a reexamination, Anal Biochem 94: 75-81, 1979.

). The KLH was estimated by a dye method using BioRad dye reagent according to the manufacture's instructions.

Conjugation of FucGM1-MMCCH to thiolated KLH

The FucGM1-MMCCH product and thiolated KLH were mixed and adjusted to pH 7.2 with 0.1M sodium phosphate buffer pH 8.0. The reaction mixture was then incubated at room temperature overnight. The content of the FucGM1-MMCCH-KLH reaction vial was transferred to a Centriprep concentrator 30 (Amicon: molecular cut-off 30000 Dalton) and unreacted FucGM1-MMCCH was removed completely with multiple washes. The conjugate was checked by HPTLC for the absence of unreacted FucGM1 as mentioned above. The epitope ratios of two batches of conjugate were determined by estimating protein content by BioRad dye binding protein assay and carbohydrate by a HPAEC-PAD assay. The epitope ratio of FucGM1-KLH (by direct method) and FucGM1-MMCCH-KLH was 149/1 and 1527/1 respectively.

B. Example 2: Synthesis of Globo-H and conjugates thereof:

1) Discussion of Synthesis:

In yet another embodiment of the present invention, an improved synthesis of Globo-H is provided utilizing the novel synthetic methodology as presented herein. The previous synthesis of globo-H described by the present inventors (Park et al. J. Am. Chem. Soc. 1996, 118, 11488; Bilodeau et al. J. Am. Chem. Soc. 1995, 117, 7840; Kim et al. J. Org. Chem. 1995, 60, 7716) utilized all glycal building blocks (Danishefsky et al.

Angew. Chem. 1996, 108, 1482; Angew. Chem. Int. Ed. Engl. 1996, 35, 1380) for the rapid construction this complex oligosaccharide. These investigations relied on a highly convergent [3+3] coupling to generate the hexasaccharide core contained in the final target. In this approach, a flexible terminal glycal was maintained throughout the hexasaccharide construction. The glycal was then used to install the ceramide side chain present en route to globo-H glycolipid 16a or its allyl glycoside 16b. The synthesis of 16a served to facilitate the proof of structure and immunocharacterization of globo-H. The allyl glycoside 16b was employed for immunoconjugation to biocarrier proteins. The previous protocols were effective in producing adequate quantities of synthetic material for proof of structure, immunocharacterization, conjugation, mouse vaccinations and phase I human clinical trials. However, improved synthetic methodologies were desired to allow for efficient bioconjugation and also to provide suitable material for clinical purposes.

10

15

20

25

30

Difficulties associated with the allyl glycoside approach invited an alternative solution which, in general terms, is described herein, and is more specifically described for fucosyl GM1 above, and additionally for Globo-H below (Figure 7). Thus, it was conceived that a hexasaccharide could be constructed containing a glycoside that would enable linkage to carrier protein, already in place (see Figure 8). Indeed this group would already have been incorporated at the reducing end of the acceptor in the [3+3] coupling step. For successful implementation of this significant new variation of the globo-H synthesis (and other complex tumor associated antigens), it would be preferable that 1) the trisaccharide acceptor containing the glycoside construct would be readily synthesizable; 2) the glycoside construct would be compatible with the [3+3] coupling; 3) the construct, in contrast to the allyl glycoside, would survive global deprotection; and 4) efficient conjugation would be implementable.

An initial retrosynthetic analysis is shown in Figure 9. For maximum convergency, the ABC acceptor was envisioned containing the aforediscussed pentenyl glycosidic linker. Additionally, the same DEF trisaccharide donor sector would be utilized as described previously. The hexasaccharide core would then be assembled via a convergent [3+3] ABC+DEF coupling reaction using a sulfonamido glycosidation protocol (Griffith et al. J. Am. Chem. Soc. 1990, 112, 5811; Griffith et al. J. Am. Chem.

Soc. 1991, 113, 5863). Previous results had indicated that the presence of a free hydroxyl at C4 of the reducing end galactose (Figure 9, see asterisk) in the DEF donor would be necessary to direct the formation of the required β-linkage in the sulfonamido glycosidation (Park et al. J. Am. Chem. Soc. 1996, 118, 11488; Kwon et al. J. Am. Chem. Soc. 1998, 120, 1588). The sequencing of the [3+3] coupling was expected to take place as shown, owing to the higher reactivity of the equatorial C3 hydroxyl group (see bold) in the acceptor as compared to the necessary axial C4 hydroxyl group in the donor trisaccharide. Important to the strategy described herein is that, once the hexasaccharide is assembled, only protecting group manipulations would be required to reach the provaccine antigen.

10

15

20

25

35

In general, the synthesis of the DEF trisaccharide sector is fairly concise, requiring six transformations starting from 6-O-TIPS galactal and tri-O-benzyl fluoro fucose (Park et al. J. Am. Chem. Soc. 1996, 118, 11488; Bilodeau et al. J. Am. Chem. Soc. 1995, 117, 7840; Kim et al. J. Org. Chem. 1995, 60, 7716). For purposes of a second-generation approach, the acceptor trisaccharide component can be dissected into a lactose derivative containing the desired NPG bearing a differentiated hydroxyl at C4' and an appropriate C-ring donor (Figure 9). The galactose donor monosaccharide also requires differential protection at C3, for eventual ABC+DEF coupling, and needs careful attention to efficiently allow for the required β-glycosidic linkage joining the AB+C domains.

As shown in Figure 10, the synthesis of the requisite ABC acceptor was conducted taking advantage of readily available lactose octaacetate, 17. Conversion of 17 to the known α-bromo donor 18 (Reithal, Y. J. Am. Chem. Soc. 1952, 74, 4210; Dasgupet et al. Carbohydr. Res. 1994, 264, 155) was followed by silver carbonate mediated glycosylation with pentenyl alcohol as acceptor, to give 19 (Pent = CH₂CH₂CH=CH₂) in 75% yield on a 100 g scale (Rodriguez et al. Aust. J. Chem. 1990, 43, 665). Similar processing of 18 with silver triflate as promoter resulted in 17% yield of the desired product. Thus, with the formation of 19, in an early stage of the synthesis, the linker was successfully installed to be used for late stage bioconjugation.

Subsequent steps were designed to generate a free acceptor site at C4' of 19 for an eventual AB + C coupling give the ABC trisaccharide (Figure 10). Removal of the ester

(

25

30

35

protecting groups in 19 to give a pentenyl lactoside was followed by a stannane mediated monobenzylation to selectively give the C3' benzyl ether (David et al. J. Chem. Soc. Perkin Trans. I 1981, 1797; Maranduba et al. Carbohydr. Res. 1986, 151, 105). In a second step, the C4' and C6' hydroxyls were engaged as a benzylidene acetal to provide compound 20 as the only observable product (Jannson et al. J. Org. Chem. 1998, 53, 5629; Koeman et al. Tetrahedron 1993, 49, 5291; Qiu et al. Liebigs Ann. 1992, 217). Finally, perbenzylation of the remaining hydroxyl groups in 20 and regioselective reductive cleavage of the benzylidene with sodium cyanoborohydride and anhydrous HCl gave the C4' alcohol 21 (Garegg, P.J. Pure Appl. Chem. 1984, 56, 845). Thus, starting from lactose octaacetate 17, the AB pentenyl glycoside acceptor 21 was obtained in 7 steps and in 20% overall yield.

With large quantities of the protected pentenyl glycoside 21 available, attention was turned to the AB+C coupling to form the trisaccharide acceptor 24. The previous synthesis of glycal 27 (Figure 10) required careful preparation of the highly activated β -fluoro donor 23 from glycal 22. The C3 PMB ether contained in 22 was strategically incorporated to allow for eventual ABC+DEF coupling upon selective deprotection of this group. In the course of this work, it was discovered that α -23 could be formed conveniently in high yield and on large scale. Accordingly, α -donor 23 was prepared from differentially protected glycal 22 by epoxidation, exposure to HF:pyridine to yield the cis fluoro-hydrin derivative and subsequent conversion of the resulting C2-hydroxyl to its benzyl ether. The anomeric α : β selectivity was demonstrated to be 10:1 and the overall yield in transforming 22 into 23 was 76%.

The effectiveness of the AB+C coupling using previously prepared β -23 and the newly prepared α -23 with the AB acceptor 21 was then investigated. The synthetic optimization of glycal trisaccharide 27 as a model case (see 23+26->27) was also examined because of its presumed sensitivity to overly demanding coupling promoters. In these investigations it was discovered that the reduced reactivity of α -fluoro donors could be attenuated by conducting the couplings with highly fluorophilic promoters in judiciously chosen solvents, as summarized in Table 1. The previous coupling procedure using the predominantly β -fluoro donor 23 and glycal 26 to give glycal trisaccharide 27 employed Muykiyama coupling conditions (Mukaiyama et al. Chem. Lett. 1981, 431;

Nicolaou et al. J. Am. Chem. Soc. 1990, 112, 3693; Nicolaou et al. J. Chem. Soc. Chem. Commun. 1991, 870) and proceeded in 54% yield with modest anomeric selectivity (entry 1, Table 1). Investigations using other promoters with α-23 are shown in entries 2 and 3, but produced little satisfaction in terms of overall efficiency. However, the preparation of glycal 27 was successfully extended to include the described α-donor 23 using strongly fluorophilic Cp₂Zr(OTf)₂ promotion (73% yield, entry 4). Gratifyingly, these optimized glycosidation conditions for formation of 27 were successfully applied to the AB+C coupling employing pentenyl glycoside 21 to provide trisaccharide 24 in yields that rivaled the parent reaction (80% yield, entry 6). Muykiyama coupling of β-23 with 21 yielded 42% of trisaccharide 24 (entry 5). Satisfied with the events leading to smooth formation of large quantities of 24, the [3+3] coupling would be investigated. The discharge of the lone PMB group in 24 could be effected in excellent yield (92%), thus completing the assembly of the desired ABC pentenyl acceptor 25.

10

15

20

25

	Acceptor AB	Donor C	Promoter, Solvent	α:β selectivity	Yield, Product
1	26ª	23 (β) ^a	SnCl ₂ , AgClO ₄ , Et ₂ O	3:1	54% (α), 18% (β) 27
2	26	23 (10α:1β)	Sn(OTf) ₂ Toluene:THF (5:1)	8:1	40% (α), 5% (β) 27
3	26	23 (10α:1β)	Cp ₂ ZrCl ₂ , AgClO ₄ , CH ₂ Cl ₂ :Et ₂ O	2.7:1	55% (α) 27
4	26	23 (10α:1β)	Cp ₂ Zr(OTf) ₂ , Toluene:THF (10:1)	10:1	72% (α), 8% (β) 27
5	21	23 (β)	SnCl ₂ , AgClO ₄ , Et ₂ O	3:1	42% (α) 24
6	21	23 (10α:1β)	Cp ₂ Zr(OTf) ₂ , Toluene:THF (5:1)	10:1	80% (α), 8% (β) 24

Table 1. Coupling conditions used to generate the ABC trisaccharide.

(a: see Park et al. J. Am. Chem. Soc. 1996, 118, 11488; Bilodeau et al. J. Am. Chem. Soc. 1995, 117, 7840; Kim et al. J. Org. Chem. 1995, 60, 7716).

The key step and final transformations completing the synthesis of 16c are shown in Figure 11. Treatment of the known DEF donor 28 (see Figure 10) with MeOTf (Lönn, H. Carbohydr. Res. 1985, 134, 105; Lönn, H. J. Carbohydr. Chem. 1987, 6, 301) In the presence of acceptor 25 smoothly provided hexasaccharide 29 in 60% yield. The configuration of the new anomeric center of 29 was confirmed to be β -configured. The [3+3] coupling yield using trisaccharide acceptor 25 was comparable to the [3+3]

procedure using the glycal-based acceptor corresponding to 27. The tremendous advantage of using 15, however, is manifested in the steps which follow.

Global deprotection began with subjection of 29 to TBAF in order to remove the silyl ethers and the cyclic carbonate. The benzyl and sulfonamido protecting groups on the resulting penta-ol were then cleaved under the action of dissolving metal reduction. This protocol was followed by peracetylation to give the isolable hexasaccharide peracetate 30. As in earlier steps, the pentenyl linkage proved highly reliable under the listed deprotection conditions. It is again notable by contrast that the corresponding allyl glycoside (to ultimately yield 16b) is not stable to the reducing metal conditions required for global deprotection and therefore must be installed subsequent to deprotection. Deacetylation of 30 with methoxide yielded the fully deprotected pentenyl glycoside of globo-H, 16c, notably poised for bioconjugation. Importantly, in the second generation variation, progress toward 16d from hexasaccharide construct 29 was greatly simplified because the need for additional functionalization to allow for conjugation is eliminated.

Toward the goal of facilitating clinical evaluation of synthetic globo-H, 16c has been conjugated to carrier protein KLH for purposes of creating a functional vaccine. The first step of this procedure involved ozonolysis of the pendant olefin, followed by reductive work-up, to give the uncharacterized aldehyde intermediate 31, as shown in Figure 12. Reductive amination with KLH and sodium cyanoborohydride in phosphate buffer yielded vaccine glycoconjugate 16d (n = 3). Covalent attachment of the carbohydrate to the proteins presumably occurs through the ε -amino groups on exposed lysine residues in KLH. Hydrolytic carbohydrate analysis of 16d revealed approximately 350 carbohydrate residues per molecule of carrier protein.

2) Experimentals:

(

10

15

20

25

30

ĺ

Peracetyl pentenyl- β -D-lactoside (19). To a cooled (ice bath) suspension of lactose octaacetate (100.0 g, 147.7 mmol), glacial acetic acid (30 mL) and acetic anhydride (30 mL) was added 100 mL of 30% HBr in AcOH dropwise over a period of 60 minutes. The reaction mixture stirred for 1 hour and the ice bath was removed. Upon stirring for an additional 2 hours at room temperature, the mixture became a homogeneous yellow solution. The solution was diluted with H_2O (1000 mL) and

extracted with CHCl₃ (3x400 mL). The organic extracts were washed with H₂O (2x1000 mL), satd. NaHCO₃ (3x500 mL), dried over MgSO₄ and concentrated. The product was azeotroped with anhydrous benzene and dried under high vacuum to yield 98.8 g (96%) of the lactosyl bromide which was used without further purification.

10

20

To a suspension of Ag₂CO₃ (100 g, 362.6 mmol), freshly activated molecular sieves (15 g) and a crystal of I₂ in 400 mL CH₂Cl₂ was added pentenyl alcohol (5.0 equiv., 73.4 mL) and then the lactosyl bromide (98.8 g, 141.4 mmol) in 400 mL of CH₂Cl₂. After stirring in the dark at room temperature for 16 hours, the reaction was filtered through a plug of Celite with additional CH2Cl2 and concentrated to a yellow oil which was purified by flash column chromatography (10% EtOAc/hexanes→50% EtOAc/hexanes) to yield 74.7 g (75%) of the pentenyl lactoside as a white foam. $[\alpha]_{22}^D$ -48.9° (c 7.5, CHCl₃); IR (film CHCl₃) 2941, 1751, 1369, 1224, 1054 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz) δ 5.60 (m, 1H), 5.17 (d, 1H, J=2.7 Hz), 5.02 (m, 1H), 4.93 (dd, 1H, J=7.9, 10.3 Hz), 4.85 (d, 1H, J=1.6 Hz), 4.78 (m, 2H), 4.71 (dd, 1H, J=9.6, 7.9 Hz), 4.30 (m, 3H), 3.93 (m, 3H), 3.66 (m, 3H), 3.45 (m, 1H), 3.30 (m, 1H), 1.98 (s, 3H), 1.94 (s, 3H), 1.91 (m, 2H), 1.89 (s, 3H), 1.88 (s, 6H, 2xCH3), 1.87 (s, 3H), 1.79 (s, 3H), 1.49 (m, 2H); ¹³C NMR (CDCl₃, 100 MHz) 8 170.33, 170.28, 170.09, 170.00, 169.74, 69.54, 169.01, 137.72, 115.00, 101.01, 100.51, 76.27, 72.76, 72.48, 71.64, 70.94, 70.58, 69.23, 69.01, 66.52, 61.97, 60.73, 29.75, 28.49, 20.80, 20.75, 20.64, 20.57, 20.45. FAB-HRMS calc'd for $C_{31}H_{44}O_{18}Na^{+}$; 727.2425. Found; 727.2418.

Pent-4-enyl 3'-O-benzyl-4', 6'-O-benzylidenyl-β-D-lactoside (20). Peracetylated pentenyl lactoside, 8, (18.2 g, 25.8 mmol) was dissolved in 300 mL of anhydrous MeOH and 2.0 mL of NaOMe (25% in MeOH) was added. The reaction stirred at rt for 16 hours and was neutralized with Dowex-H⁺ (pH 5-6). The reaction was filtered with additional MeOH and concentrated to a white solid, 19a, (10.6 g, quantitative) which was used without further purification: ¹H NMR (D₂O, 400 MHz) δ 5.81 (m, 1H), 5.00 (dd, 1H, J=17.3, 1.9 Hz), 4.92 (dd, 1H, J=8.9 Hz), 4.74 (m, 1H), 4.39 (d, 1H, J=8.0 Hz), 4.35 (d, 1H, J=7.8 Hz), 3.72-3.42 (m, 12H), 3.21 (m, 1H0, 2.06 (m, 2H), 1.63 (m, 2H); ¹³C NMR (D₂O, 100 MHz) δ 141.27, 117.31, 105.42, 104.54, 80.85, 77.84, 77.24, 76.92, 75.33, 75.00, 73.44, 72.47, 71.03, 63.52, 62.56, 31.83, 30.48.

5

10

15

20

25

30

The hepta-ol 19a (1.14 g, 2.8 mmol) and dibutyltin oxide (0.76 g, 3.1 mmol) were heated at reflux in benzene (70 mL) with azeotropic water removal for 15 h. The mixture was doubled in concentration, cooled to room temperature, and benzyl bromide (0.69 ml, 5.8 mmol) and Bu₄NI (1.03 g, 2.8 mmol) were added. The mixture was heated at reflux 3.5 h, cooled, silica gel was added to the flask, and the solvent was evaporated. The residue was applied to a column of silica gel, tin by-products were removed by flushing with hexanes, and elution (5 % MeOH in CH₂Cl₂) gave the pure 3'-O-benzyl ether (0.76 g, 54 %) as a white foam: $[\alpha]_{22}^D + 36.7^\circ$ (c 2.73, CHCl₃); IR (film CHCl₃) 3371, 2924, 2880, 1372, 1157, 1074 cm⁻¹; ¹H NMR (MeOH-d₄, 400 MHz) δ 7.46-7.40 (m, 2H), 7.35-7.20 (m. 3H), 5.92-5.72 (m, 1H), 5.08-4.93 (m, 2H), 4.76 (d, 1H, J=11.8 Hz), 4.65 (d, 1H, J=11.8 Hz), 4.38 (d, 1H, J=7.8 Hz), 4.28 (d, 1H, J=7.8 Hz), 4.02 (d, 1H, J=2.9 Hz), 3.95-3.63 (m, 6H), 3.61-3.48 (m, 4H), 3.43-3.20 (m, 3H), 2.20-2.10 (m, 2H), 1.78-1.65 (m, 2H); 13 C NMR (MeOH- d_4 , 100 MHz) δ 139.77, 139.47, 129.29, 129.08, 128.64, 115.19, 105.02, 104.23, 82.17, 80.74, 76.88, 76.40, 76.35, 74.71, 72.55, 71.81, 70.23, 67.02, 62.44, 61.91, 31.22, 30.07. FAB-HRMS calc'd for C₂₄H₃₆O₁₁Na⁺; 523.2155. Found; 523.2137.

The 3'-O-benzyl ether (0.6 g, 1.20 mmol) was dissolved in acetonitrile and DMF (5:2, 7 mL), and benzaldehyde dimethylacetal (0.47 mL, 3.1 mmol) and CSA (14 mg, 60 μ mol) were added. After stirring 16 h at room temperature, the mixture was diluted with CH₂Cl₂ and washed with sat. NaHCO₃. The organic extracts were dried (MgSO₄), evaporated, and following addition of ether (100 mL) to the resulting residue, the pure 20 was recovered by filtration (0.51 g, 72 %): $[\alpha]_{22}^D +111^\circ$ (c 2.21, MeOH); IR (CHCl₃ film) 3440, 2872, 1368, 1163, 1109, 1048, 1005 cm⁻¹; ¹H NMR (MeOH- d_4 , 400 MHz) δ 7.55-7.11 (m, 10H), 5.82-5.69 (m, 1H), 5.45 (s, 1H), 4.98-4.83 (m, 2H), 4.64 (d, 2H, J=3.0 Hz), 4.40 (d, 1H, J=7.9 Hz), 4.23 (d, 1H, J=3.4 Hz), 4.18 (d, 1H, J=7.8 Hz), 4.15-3.98 (m, 2H), 3.87-3.66 (m, 4H), 3.55-3.10 (m, 7H), 2.20-2.10 (m, 2H), 1.65-1.53 (m, 2H); ¹³C NMR (MeOH- d_4 , 100 MHz) δ 139.76, 139.49, 139.47, 129.86, 129.30, 129.07, 129.03, 128.72, 127.35, 115.19, 104.69, 104.28, 102.03, 80.63, 80.17, 76.37, 76.28, 74.77, 74.73, 72.84, 70.86, 70.25, 68.17, 61.70, 31.22, 30.07. FAB-HRMS calc'd for $C_{31}H_{40}O_{11}Na^+$; 611.2468. Found; 611.2465.

Pent-4-enyl 2, 2', 3, 3', 6, 6'-hexa-O-benzyl-β-D-lactoside (21). The tetraol 20 (0.51 g, 0.87 mmol) and Et₄NI (0.12 g, 0.43 mmol) were dried (azeotropic distillation with benzene), dissolved in DMF (5 mL) and cooled to 0°C. Benzyl bromide (0.83 mL, 7.0 mmol) was added followed by NaH (0.22 g, 60 %, 5.6 mmol) and the mixture was allowed to warm to room temperature over 14 h. The mixture was diluted with ethyl acetate, washed with water, the organic layer was dried (MgSO₄) and evaporated. 10 Purifiction of the residue by chromatography on silica gel (4:1->2:1 hexanes:ethyl acetate) gave pure pentabenzyl lactoside as a white foam (0.80 g, 97 %): $[\alpha]_{22}^D$ +129° (c 1.63, CHCl₃); IR (CHCl₃ film) 3030, 2866, 1453, 1365, 1096, 1063, 1028, 911 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz) δ 7.50-7.05 (m, 30H), 5.80-5.65 (m, 1H), 5.38 (s, 1H), 5.10 (d, 1H, J=10.6 Hz), 4.99-4.60 (m, 9H), 4.47 (d, 1H, J=12.1 Hz), 4.38 (d, 1H, J=7.8 Hz), 4.30 (d, 1H, J=7.8 Hz), 4.25 (d, 1H, J=12.1 Hz), 4.12 (d, 1H, J=13 Hz), 3.94 (d, 1H, J=3.4 Hz), 3.92-3.60 (m, 6H), 3.54 (dd, 1H, J=8.8 Hz, 9.2 Hz), 3.46 (dd, 1H, J=2.6 Hz, 7.0 Hz), 3.40-3.23 (m, 3H), 2.85 (s, 1H), 2.22-2.00 (m, 2H), 1.75-1.60 (m, 2H); ¹³C NMR (CDCl₃, 100 MHz) δ 138.92, 138.63, 138.51, 138.04, 128.80, 128.52, 128.31, 128.24, 128.17, 128.13, 128.06, 128.03, 128.00, 127.69, 127.65, 127.54, 127.49, 127.38, 127.30, 126.52, 20 114.84, 103.59, 102.83, 101.30, 83.01, 81.81, 79.60, 78.76, 77.65, 75.73, 75.22, 75.05, 74.97, 73.61, 72.91, 71.56, 69.27, 68.90, 68.27, 66.28, 30.18, 28.89. FAB-HRMS calc'd for C₅₉H₆₄O₁₁Na⁺; 971.4346. Found; 971.4375. The benzylidene (0.63 g, 0.66 mmol) was dissolved in THF (6.6 mL) and stirred with

The benzylidene (0.63 g, 0.66 mmol) was dissolved in THF (6.6 mL) and stirred with freshly activated 4ÅMS (0.25 g) 10 min at room temperature. In one portion NaCNBH₃ (0.21 g, 3.3 mmol) was added followed by anhydrous HCl (2.0 M Et₂O), dropwise until the mixture no longer bubbled (approx. 2 mL). After stirring and additional 10 min, the mixture was passed through a plug of Celite washing with ethyl acetate, the filtrate was washed with sat. NaHCO₃ and brine, dried (MgSO₄) and the organic layers evaporated. Purification by column chromatography (9:1 hexanes:ethyl acetate) gave pure 21 as white solid (0.49 g, 79 %): $[\alpha]_{22}^D$ +200° (c 1.05, CHCl₃); IR (CHCl₃ film) 3474, 3062, 3029, 2869, 1453, 1364, 1094, 1028 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz) δ 7.40-7.06 (m, 30H), 5.80-5.66 (m, 1H), 5.02-4.85 (m, 3H), 4.81 (d, 1H, J=11.0 Hz), 4.75-4.54 (m, 6H), 4.67 (d, 1H, J=12.2 Hz), 4.42-4.26 (m, 5H), 3.94 (s, 1H), 3.92-3.81 (m, 2H), 3.71 (dd, 1H, J=10.7 Hz, 4.1 Hz), 3.64 (d, 1H, J=10.6 Hz), 3.57 (dd, 1H, J=9.4 Hz, 5.5 Hz), 3.55-3.42

25

35

1

(m, 3H), 3.38 (dd, 1H, J=5.2 Hz, 9.6 Hz), 3.36-3.21 (m, 4H), 2.32 (s, 1H), 2.15-2.02 (m, 2H), 1.74-1.60 (m, 2H); ¹³C NMR (CDCl₃, 100 MHz) δ 139.04, 138.54, 138.52, 138.23, 138.09, 137.96, 137.81, 128.33, 128.15, 127.93, 127.66, 127.50, 114.80, 103.50, 102.43, 82.79, 81.68, 80.99, 79.27, 76.52, 75.22, 75.10, 74.99, 74.83, 73.37, 72.99, 72.67, 71.86, 69.10, 68.32, 68.16, 66.00, 30.11, 28.83. FAB-HRMS calc'd for C₅₉H₆₆O₁₁Na⁺; 973.4503. Found; 973.4513. α-Fluoro donor (23). A solution of 3-O-PMB-4,6-Di-O-benzyl-galactal (2.24 g, 5.02 mmol) in dry CH₂Cl₂ (5 ml) under N₂ at 0°C was treated with dimethyldioxirane (0.11 M, 47 ml), and the mixture was stirred until all of the glycal was consumed (~ 1h, TLC 30% EtOAc in hexane) Note: Elevated temperature and/or excess of DMDO will prompt oxidation of the PMB group and lower reaction yield. The solvents were evaporated under vacuum at 0°C and the residue was kept under high vacuum. The flask containing galactal epoxide was charged with freshly prepared 4 A molecular sieves (2 g), dry THF (50 ml) and cooled to 0°C. HF/Pyr complex (0.79 ml, ~5 equiv.) was added dropwise via syringe. The reaction mixture was left overnight to slowly reach room temperature and quenched with Et₃N (1.27 g, \sim 2.5 equiv.) to reach pH \sim 7. The mixture was filtered 20 through a pad of anhydrous MgSO4 and rinsed three times with 50 ml of EtOAc. The filtrate was washed with water (50 ml) and saturated NaHCO3 solution (50 ml), dried over MgS04 and concentrated to dryness. Flash column chromatography (EtOAc/hexanes, 2/1) gave 2.06 g (85% yield) of fluorohydrin as a mixture of anomers α : $\beta = 10$: 1. ¹⁹F NMR (CDCl₃, 376 MHz, C₆F₆ as external standard) δ 9.7 (dd, α , J = 54.4, 25.0 Hz) 20.0 (dd, β , J = 53.9, 13.1 Hz); ¹H NMR (CDCl₃, 400 MHz) δ 7.38 – 7.24 (m, 12H), 6.90 (d, 2H, J = 8.7 Hz), 5.70 (dd, 1H, J = 54.4, 2.8 Hz), 4.89 and 4.57 (two d, 2H, J = 11.3 Hz), 4.70 and 4.54 (AB q, 2H, J = 11.2 Hz), 4.54 and 4.46 (AB q, 2H, J = 11.8 Hz), 4.17 (AMX octet, 1H, J = 2.8, 10.1, 25.0 Hz), 4.13 (br t, 1H, J = 6.8Hz), 4.06 (d, 1H, J = 1.5 Hz), 3.81 (s, 3H), 3.74 (dd, 1H, J = 2.6, 10.1 Hz), 3.60 (m, 2H). The above mixture (8.29 g, 17.2 mmol) was dissolved in dry DMF (100 ml) containing freshly prepared 4 A molecular sieves (3 g) under N₂ at 0°C, treated with benzyl bromide (4.41 g, 25.8 mmol, 1.5 equiv.) and finally with NaH (1.24 g, 60% dispersion in oil, 30.86 mmol, 1.8 equiv.), and stirred overnight at room temperature. The reaction was

quenched with glacial acetic acid (0.93 g, 0.9 equiv.) and the mixture filtered through a

1

10

20

25

30

pad of anhydrous MgSO₄ with EtOAc (4x50 ml). The organic solution was washed with water (4 x 50 ml), dried (MgSO₄) and concentrated in vacuo. chromatography of the residue (hexane/EtOAc, 4/1) gave 9.36 g (95%) of the title compound as colorless liquid with the same ratio of anomers $\alpha:\beta=10:1$ as the starting fluorohydrin. ¹⁹F NMR (CDCl₃, 376 MHz, C_6F_6 as external standard) δ 11.5 (dd, α , J =53.7, 25.2 Hz), 22.8 (dd, β , J = 53.4, 13.0 Hz). For analytical purpose 50 mg of pure α anomer was obtained using preparative HPLC. [α]₂₂D - 54.5° (c 0.55, CHCl₃); ¹H NMR (CDCl₃, 500 MHz), δ 7.38–7.24 (m, 17H), 6.88 (d, 2H, J=8.6 Hz), 5.58 (dd, 1H, J=53.7, 2.7 Hz, 4.93 (d, 2H, J=11.34 Hz), 4.56 (d, 2H, J=11.34 Hz), 4.85 (AB q, 2H, 2Hz), 4.85 (AB q, 2Hz), 4.85 (AB q, 2Hz), 4.85 (AB q, 2Hz) 11.78 Hz), 4.72 (AB q, 2H, J = 11.78 Hz), 4.73 (AB q, 2H, J = 11.3 Hz), 4.68 (AB q, 2H, J = 11.3 Hz), 4.47 (AB q, 2H, J = 11.84 Hz), 4.41 (AB q, 2H, J = 11.84 Hz), 4.09 (br t, 1H, J = 6.5 Hz), 4.02 (AMX m, 1H, J = 2.7, 10.05, 25.2 Hz), 3.98 (app s, 1H), 3.92 (dd, 1H, J = 2.64, 10.05 Hz), 3.81 (s, 3H), 3.54 and 3.52 (ABX m, 2H, J=9.3, 6.05, 7.0 Hz); ¹³C **NMR** (CDCl₃, 125MHz) 8 159.20, 138.35, 138.08, 137.71, 130.43, 129.18, 128.39, 128.25, 128.14, 127.92, 127.8, 127.78, 127.66, 113.81, 106.25 (d, J = 229.0 Hz), 78.09, 75.65 (d, J = 23.5 Hz), 74.79(ArCH₂), 74.29, 73.70 (ArCH₂), 73.45 (ArCH₂), 72.71 (ArCH₂), 71.70 (d, J = 2.7 Hz)68.26.55.24 (CH₃O); MS (NH₃) 586 ([M + NH₄]⁺). PMB trisaccharide (24). A mixture of lactoside (21) (402 mg, 0.423 mmol) and fluoro donor (23) (485 mg, 0.846 mmol, 2 equiv.) was azeotroped with anhydrous benzene (3x10 ml) and further dried on high vacuum for 3 h. The above mixture was dissolved in toluene (3.8 ml) and transferred via cannula to a flask containing freshly prepared 4 Å molecular sieves (0.68 g) under N2, treated with 2,6-di-tert-butylpyridine (143 µl) and cooled to -20 °C. (Cp)₂Zr(OTf)₂ (225 mg, 0.381 mmol, 0.9 equiv.) was suspended in THF (0.38 ml) and added via a cannula to the reaction mixture. The reaction was stirred for 72 h at 7 °C in darkness. The reaction mixture was diluted with EtOAc (10 ml) and filtered through a pad of anhydrous MgSO₄ with EtOAc (3x 10mL). The filtrate was washed with satd. NaHCO3 solution (2 x 10 ml), dried over MgSO4, and concentrated to dryness. Flash column chromatography (2% Et₂O/CH₂Cl₂) gave 509 mg (80%) of desired α -product (24) and 51 mg (8%) of β -product. $[\alpha]_{22}^D$ +24.6° (c 3.90, CHCl₃); IR (CHCl₃ film) 3062, 3029, 2919, 2868, 1612, 1513, 1496, 1364, 1303, 1248, 35

5 1028 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz) δ 7.40-6.95 (m, 49 H), 6.69 (d, 1H, J=8.5 Hz), 5.73 (m, 1H), 5.00-4.93 (m, 2H), 4.92-4.84 (m, 2H), 4.82-4.73 (m, 2H), 4.72-4.63 (m, 5H), 4.61 (d, 1H, J=13.0 Hz), 4.48-4.35 (m, 5H), 4.34-4.24 (m, 4H), 4.16 (d, 2H, J=6.8 Hz), 4.07 (dd, 1H, J=8.8 Hz), 4.02-3.80 (m, 8H), 3.78-3.60 (m, 3H), 3.68 (s, 3H), 3.60-3.35 (m, 6H), 3.35-3.18 (m, 4H), 3.12-3.04 (m, 1H), 2.06 (m, 2H), 1.65 (m, 2H); ¹³C NMR (CDCl₃, 100 MHz) δ 158.76, 139.66, 139.45, 139.26, 139.16, 139.09, 138.92, 138.57, 138.52, 131.39, 129.30, 128.95, 128.70, 128.60, 129.30, 128.08, 127.95, 115.35, 114.02, 104.05, 103.35, 101.25, 83.14, 82.17, 79.91, 79.71, 77.77, 77.04, 75.69, 75.58, 75.46, 75.33, 74.17, 73.75, 73.54, 73.48, 72.65, 72.54, 69.91, 69.71, 68.80, 68.33, 68.19, 55.11, 30.14, 28.86; FAB-HRMS calc'd for C₉₄H₁₀₂O₁₇Na⁺; 1525.7014. Found; 1525.6996.

Trisaccharide acceptor (25). A solution of PMB trisaccharide (24) (445 mg, 0.296 mmol) in methylene chloride (10 ml) at 0 °C was treated with phosphate buffer (1.5 ml, pH = 7.4) and DDQ (89 mg, 1.3 equiv.) and stirred at 0 °C for 5 h. The reaction mixture was diluted with EtOAc (50 ml), washed with satd. NaHCO₃ solution (2 x 20 ml) and water (20 ml), dried over MgSO₄, and concentrated to dryness. The crude material was purified by flash column chromatography (4% ether in methylene chloride) to give 344 mg (84%) of deprotected trisaccharide (25) as a colorless oil. $[\alpha]_{22}^D$ +28.2° (c 5.70, CHCl₃); IR (CHCl₃ film) 3570, 3062, 3029, 2913, 2868, 1496, 1453, 1364, 1208, 1095 cm⁻¹: ¹H NMR (CDCl₃, 400 MHz) δ 7.77-7.06 (m, 45H), 5.73 (m, 1H), 5.01 (dd, 1H, J=5.5, 3.3 Hz), 4.95 (dd, 1H, J=5.8, 2.6 Hz), 4.90 (m, 1H), 4.78 (d, 1H, J=10.9 Hz), 4.75 (d, 1H, J=11.4 Hz), 4.70-4.59 (6d, 6H), 4.47-4.37 (m, 5H), 4.28 (m, 3H), 4.19 (s, 2H), 4.08-3.91 (m, 6H), 3.85 (m, 2H), 3.69 (m, 5H), 3.66 (1H, d, J=11.0 Hz), 3.50-3.19 (m, 9H), 3.10 (dd, 1H), 2.07 (m, 2H), 1.79 (d, 1H, OH), 1.65 (d, 2H); ¹³C NMR (CDCl₃, 100 MHz) & 139.36, 138.72, 138.63, 138.52, 138.41, 138.29, 138.19, 138.07, 137.98, 128.35, 128.20, 128.06, 127.97, 127.66, 127.54, 127.08, 114.82, 103.55, 102.67, 99.58, 82.93, 81.67, 81.55, 79.32, 77.61, 76.90, 75.13, 75.02, 74.96, 74.80, 73.08, 72.99, 72.91, 72.01, 69.95, 69.22, 69.15, 68.34, 67.73, 67.57, 60.35, 30.19, 28.92; FAB-HRMS calc'd for C₈₆H₉₄O₁₆Na⁺; 1405.6439. Found; 1405.6385.

20

25

30

35

Hexasaccharide (29). The thioethyl donor 28 (543 mg, 0.425 mmol) and acceptor 25 (587 mg, 0.425 mmol) were combined, azeotroped with anhydrous benzene (5x5 mL)

10

20

25

30

and placed under high vacuum for 5 hours. The mixture was then dissolved in 3.5 mL CH2Cl2 and 7.0 mL Et2O, treated with freshly prepared molecular sieves and cooled to 0 °C. Methyl triflate (3.0 equiv., 144 µL) was added in one portion and the reaction stirred at 0 °C for 3 hours. Another 144 µL of MeOTf was added and the reaction was allowed to stir for an additional 2 hours at 5 °C. The reaction was quenched by the addition of solid NaHCO3, filtered through Celite with EtOAc, concentrated and purified by HPLC (17% EtOAc/hexanes) to give 663 mg (60%) of hexasaccharide as a white foam. $[\alpha]_{22}^D$ – 9.7° (c 1.00, CHCl₃); IR (CHCl₃ film) 3533, 3343, 3087, 3030, 2940, 2865, 1790, 1496, 1453, 133, 1095 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz) δ 7.76 (d, 2H, J=7.5 Hz), 7.45-7.00 (m, 63H), 5.84 (m, 1H), 5.20 (s, 1H), 5.11 (d, 1H, J=3.2 Hz), 5.09 (d, 1H, J=3.6 Hz), 5.05 (d, 1H, J=3.3 Hz), 5.03 (m, 1H), 4.92 (m, 2H), 4.86 (d, 1H, J=6.0 Hz), 4.82 (m, 2H), 4.78 (1H, d, J=2.2 Hz), 4.74-4.61 (m, 8H), 4.53-4.44 (4d, 4H), 4.38-4.30 (m, 4H), 4.18-3.82 (m, 20H), 3.76-3.66 (m, 5H), 3.66-3.60 (m, 2H), 3.58-3.52 (m, 2H), 3.48-3.40 (m, 2H), 3.38-3.32 (m, 2H), 3.29-3.25 (m, 3H), 3.06 (dd, 1H, J=10.2 Hz), 2.86 (s, 1H), 2.74 (m, 1H), 2.16 (m, 2H), 1.74 (m, 2H), 1.23 (s, 3H, J=6.5 Hz), 1.16-1.07 (m, 42H); ¹³C NMR (CDCl₃, 100 MHz) & 155.49, 140.71, 139.37, 138.96, 138.72, 137.70, 138.66, 138.55, 138.42, 138.37, 138.10, 138.07, 138.04, 137.88, 132.07, 128.89, 128.64, 128.50, 128.27, 128.16, 128.04, 127.86, 127.68, 127.53, 127.34, 127.20, 114.79, 103.49, 103.14, 102.61, 99.63, 99.12, 97.79, 82.26, 81.61, 81.34, 80.45, 79.36, 78.95, 78.26, 77.82, 77.64, 77.45, 77.24, 77.16, 76.83, 76.45, 75.39, 75.28, 75.12, 74.98, 74.89, 74.78, 73.94, 73.13, 72.94, 72.92, 72.52, 71.91, 71.81, 71.25, 71.11, 69.35, 69.23, 69.18, 68.18, 68.11, 68.01, 67.77, 67.54, 61.98, 61.72, 56.03, 30.16, 28.88, 18.01, 18.00, 17.95, 17.92, 11.85, 11.82; LRMS (FAB) calc'd for C₁₅₀H₁₈₅NO₃₂SSi₂Na⁺ 2624.

Peracetate of globo-H pentenyl glycoside (30). To a solution of the hexasaccharide (585 mg, 0.224 mmol) in THF (10 mL) was added TBAF (1.0 M THF, 10 equiv., 2.24 mL). The reaction stirred at rt for 3 days, poured into ice water and extracted with EtOAc (3x50 mL). The organic extracts were washed with satd. NaHCO₃ (50 mL) and brine (50 mL), dried over MgSO₄ and concentrated to an oil which was purified through a short plug of silica gel with EtOAc. The resulting triol was dissolved in anhydrous MeOH (8 mL) and sodium methoxide was added (0.25 mL of a 25% solution in MeOH).

The reaction stirred at rt for 18 hours, neutralized with Dowex-H+, filtered with MeOH washings and concentrated. To the resulting white solid was added THF (2.0 mL) and condensed liquid NH3 (~25 mL) at -78 °C. Sodium (~500 mg) was added and the resulting blue solution stirred at -78 °C for 2 hours. The reaction was quenched with anhydrous MeOH (~10 mL), brought to rt and concentrated under a stream of dry N2 to a volume of ~5 mL. The reaction was neutralized with Dowex-H+, filtered with MeOH 10 washing and concentrated to a white solid. The white solid was dissolved in 5.0 mL pyridine and 5.0 mL CH2Cl2 and cooled to 0 °C. A crystal of DMAP was added followed by acetic anhydride (5.0 mL). The ice bath was removed and the reaction stirred at rt overnight. Concentration followed by purification by flash column (gradient elution 75% EtOAc/hexanes→100% chromatography 15 MeOH/EtOAc) gave 168 mg (42%) of 30 as a white solid: $[\alpha]_{22}^D$ 4.37° (c 1.85, CHCl₃); IR (CHCl₃ film) 2939, 1747, 1370, 1229, 1066 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz) δ 7.66 (d, 1H, J=6.5 Hz), 5.77 (m, 1H), 5.58 (d, 1H, J=3.2 Hz), 5.47 (d, 1H, J=3.5 Hz), 5.39 (d, 1H, J=3.2 Hz), 5.29 (dd, 1H, J=10.9, 3.0 Hz), 5.24-5.06 (m, 5H), 5.04-5.02 (m, 1H), 4.99-4.85 (m, 4H), 4.74 (dd, 1H, J=10.9, 2.9 Hz), 4.53-4.40 (m, 5H), 4.36 (m, 1H), 4.26 20 (dd, 1H, J=10.6, 3.4 Hz), 4.18-4.03 (m, 6H), 3.99-3.96 (m, 2H), 3.87-3.81 (m, 3H), 3.77-3.74 (m, 1H), 3.51-3.45 (m, 1H), 3.03 (m, 1H), 2.16 (s, 3H), 2.15 (s, 3x3H), 2.13-2.11 (m, 2H), 2.10 (s, 3H), 2.09 (s, 3H), 2.08 (s, 2x3H), 2.08 (s, 3H), 2.07 (s, 3H), 2.05 (s, 3H), 2.04 (s, 2x3H), 2.00 (s, 3H), 1.97 (s, 2x3H), 1.89 (s, 3H), 1.65 (m, 2H), 1.62 (s, 3H), 1.14 (d, 3H, J=6.5 Hz); ¹³C NMR (CDCl₃, 100 MHz) δ 172.31, 171.55, 170.78, 170.61, 25 170.57, 170.48, 170.41, 170.30, 170.08; 169.75, 169.61, 169.57, 169.44, 168.96, 137.76, 115.07, 102.05, 101.29, 100.45, 99.23, 98.74, 94.29, 77.24, 77.16, 76.07, 73.68, 73.40, 73.17, 72.63, 72.34, 71.85, 71.77, 71.56, 71.34, 70.83, 70.71, 70.19, 70.08, 69.32, 69.03, 68.88, 68.09, 68.01, 67.59, 67.32, 64.48, 29.80, 28.54, 23.12, 20.90, 20.88, 20.82, 20.74, 20.73, 20.72, 20.71, 20.64, 20.62, 20.55, 20.54, 20.49, 15.91; FAB-HRMS calc'd for 30 C₇₇H₁₀₇NO₄₇Na⁺;1820.5911. Found; 1820.5994.

Globo-H, pentenyl glycoside (16c). The peracetate (20 mg, 0.011 mmol) was dissolved in anhydrous MeOH (2.0 mL) and 100 µL of sodium methoxide was added (25% solution in MeOH). The reaction stirred at rt for 18 hours, was neutralized with

1

Dowex-H⁺ (-pH 6-7), filtered with MeOH washings, concentrated and purificated using RP silica gel (H₂O →1% MeOH/H₂O) then P-2 Gel (H₂O elutent) to yield 12 mg (99%) of a white solid. [α]₂₂^D 3.00° (c 1.00, MeOH); IR 3374, 2930, 1641, 1372, 1070 cm⁻¹; ¹H NMR (MeOH, 400 MHz) δ 5.79 (m, 1H), 5.18 (d, 1H, J=3.9 Hz), 4.98 (dm, 1H, J=7.2 Hz), 4.91 (m, 1H), 4.87 (m, 1H), 4.51 (s, 1H), 4.49 (d, 1H, J=1.4 Hz), 4.41-4.36 (m, 2H), 4.24-4.20 (m, 4H), 4.10 (d, 1H, J=2.5 Hz), 4.06-4.00 (m, 3H), 3.94 (s, 1H), 3.87-3.45 (m, 22H), 3.35-3.31 (m, 2H), 3.19 (t, 1H, J=8.8 Hz), 2.10 (m, 2H), 1.96 (s, 3H), 1.66 (m, 2H), 1.19 (d, 3H, J=6.5 Hz); ¹³C NMR (CDCl₃, 100 MHz) δ 174.53, 139.53, 115.27, 105.50, 105.44, 104.30, 103.96, 102.81, 101.07, 81.29, 80.59, 80.04, 79.16, 78.00, 76.81, 76.57, 76.49, 76.45, 76.39, 75.57, 74.89, 74.69, 73.58, 72.64, 72.49, 71.56, 70.65, 70.63, 70.38, 70.31, 69.70, 68.13, 62.63, 62.59, 61.94, 61.62, 53.11, 49.90, 31.29, 30.14, 23.55, 16.76. FAB-HRMS calc'd for C₄₃H₇₃NO₃₀Na⁺; 1106.4115. Found; 1106.4105.

C. Example 3: Preparation of glycoamino acids and inventive glycopeptides:

1) Discussion of synthetic methods:

20

35

In general, it was desired to incorporate the two antigens described above, fucosyl GM1 and globo-H into glycopeptides. As shown in Figure 13, a tranformation utilizing catalytic asymmetric hydrogenation of glycosylated amide esters was contemplated. The new approach anticipated a Homer-Emmons olefination of the protected aldehyde with a suitably protected glycine derived phosphonate to give an enamide ester. Subsequent catalytic asymmetric hydrogenation would hopefully yield diastereromerically pure glycoamino acid.

In but one example, an inventive glycoamino acid, based on a peraceylated lactose derivatives was prepared. Specifically the required lactose derived enamide ester substrate was prepared according to Figure 14. Ozonolysis of the NPG 32 (Allen et al., Chem. Eur. J. 2000, 6, 1366) followed by reductive work-up gave the corresponding aldehyde derivative. The crude aldehyde was then subjected to Horner-Emmons olefination using tetramethylguanidine and phosphonate 33. Phosphonate 33, with N-Boc and 2-(trimethylsilyl)ethyl ester (TSE) protection (Schmidt et al., Synthesis 1984, 53; Kawai et al., Chem. Lett. 1990, 577) was chosen because of the need for the resulting glycoamino

WO 01/14395 PCT/US00/22894

acids to be orthogonally suitable for peptide couplings in the presence of acetate carbohydrate protecting groups. The enamide ester 34 was obtained as a single geometric isomer in 88% yield for the 2-step procedure.

10

20

25

30

35

In one preferred embodiment, conditions for asymmetric hydrogenation of enamide ester 34 are detailed. The (S, S) ligand isomer of ethyl DuPHOS catalyst precursor was utilized, which has been well characterized in these types of systems to give the (S)-isomer in the amino acid product. The protected glycoamino acid was obtained in 98% yield and was determined to have been formed with a diastereomeric ratio (dr) of >20:1. Remarkably, the t-Boc protons are nearly baseline resolved and, in the asymmetric reaction, the minor isomer could not be detected. ¹³C analysis also supports the conclusion that the minor isomer is not formed within the limit of NMR detection. Hydrogenation of 34 with an achiral catalyst (Pd/C, MeOH) produced a 1:1 mixture of R and S configured 35, providing a comparison for diastereomeric ratio determination. This reaction also indicates that chirality transfer to yield 35 occurs from the chiral ligand and not carbohydrate derived substrate control. A final step to be performed prior to moving to synthesis and assembly of tumor antigens was that of demonstrating deprotectability of the blocking groups contained in the amino acid side chain. In the event, reaction of 35 with TBAF in THF gave acid 36, suitably prepared for peptide coupling, in 93% yield.

With the general methodology demonstrated in the lactose model, in other preferred embodiments, advanced hexasaccharides 37 and 38, as well as other antigens of interest, were investigated. As shown in Table 2, olefination of the peracetylated *n*-pentenyl glycoside of Globo-H, 37, under the same conditions as those used in Figure 14, provided the corresponding enamide ester 41 in 72% yield as a single isomer and provided fucosyl GM1 hexasaccharide 16 in 10-22% yield. Notably, through use of the (S, S)-Et-DuPHOS-Rh⁺ catalyst system, the hydrogenation of 41 and 42 proceeded in excellent yield producing 45 and 46 as single diastereomers by ¹H NMR analysis. Compounds 45 and 46 represent the first examples of synthetic glycoamino acids containing the complex oligosaccharides Globo-H and fucosyl GM₁.

Similar transformations on two other clinically promising antigens to generate their corresponding glycoamino acids were also performed in certain other embodiments.

As discussed previously, Lewis (Le) oligosaccharide has been identified as an important antigen for eliciting antibodies against colon, liver, prostate and ovarian carcinomas (Lloyd et al., Am. J. Clin. Path. 1987, 87, 129; Lloyd et al., Cancer Biol. 1991, 2, 421; Yin et al., Int. J. Cancer, 1996, 65, 406). Previously, both a Le -KLH conjugate vaccine (Danishefsky et al. J. Am. Chem. Soc. 1995, 117, 5701) and a clustered Le glycopeptide (of natural α-O-linked configuration) glycoconjugate attached to either a glycolipid or KLH have been prepared, and have initiated human clinical trials against ovarian cancer with these vaccines have been initiated (Kudryashov et al., Cancer Immunol. Immunother. 1998, 45, 281; Sabbatini et al., Int. J. Cancer 2000, 87, 79).

The results starting with both Le^y n-pentenyl glycoside 39 and the α-linked n-pentenyl glycoside of the Tn antigen 40 (GalNAc) are presented in Table 2. The pentasaccharide 39 was available as an intermediate in the synthesis of Le^y glycopeptide cluster and consequently illustrates the potential advantage of this strategy. Thus, if immunogenicity is retained in the artificial constructs, these NPG derived glycoamino acids offer a much shorter synthetic route to vaccine glycoconjugates than their native counterparts. As shown in Table 2, olefination of 39 and 40 was uneventful and enamide esters 43 and 44 were obtained in 85% and 75% yields respectively, again as single isomers. Asymmetric hydrogenation, as before, also produced diastereomerically pure glycoamino acids 47 and 48 in excellent yields.

10

15

Table 2

With the glycoamino acids as described above in hand, it would thus be desirable to generate novel glycopeptides. Specifically, in one embodiment, a novel glycopeptide incorporating globo-H, Le^y, and Tn is provided. Specifically, the C-terminus is modified to include a conjugation handle for carrier protein KLH. The mercaptoacetamide unit has proven to be effective for this purpose. As shown in Figure 15, the Tn glycoamino acid 48 was treated with TBAF to reveal the corresponding carboxylic acid. Coupling with a di-amino spacer terminated in a protected mercaptoacetamide (AcSCH₂C(O)(CH₂)₃NH₂) under the agency of the BOP reagent (benzotriazol-1-

WO 01/14395 PCT/US00/22894

(

10

20

25

oxytris(dimethylamino)phosphonium hexafluorophosphate) gave the corresponding amide in 50% yield for the 2 steps. Removal of the *N*-terminal Boc group gave amine 49 as its trifluoroacetate salt. The next antigen, Le^y, was prepared for coupling by reaction of 47 with TBAF to give acid 50. Coupling of amine 49 with Le^y acid 50, again with the BOP promoter, gave the Tn-Le^y di-peptide 51 in 86% yield. Lastly, Globo-H glycoamino acid 45 was treated with TBAF to give its corresponding acid 52. Removal of the Boc protecting group in 51 followed by coupling with acid 52 gave the Tn-Le^y-Globo-H tripeptide in 64% yield. Finally, the *N*-terminal Boc group was removed and the resulting anime capped as its acetate to give tripeptide 53 in 95% yield. With all components in place, the ester protecting groups were removed with hydrazine in degassed methanol to give the fully deprotected glycopeptide 54 (Figure 16) in excellent yield. As discussed below, the inventive glycopeptides prepared as detailed herein can also be conjugated to a suitable carrier protein or lipid.

2) Experimental General. DuPHOS-Rh⁺ catalysts were purchased from Strem Chemical Co., Newburyport, MA. All other commercial materials (purchased from Aldrich-Sigma) were used without further purification. The following solvents were obtained from a dry solvent system (passed through a column of alumina): THF, diethyl ether (Et₂O), CH₂Cl₂, toluene and benzene. All reactions were performed under an atmosphere of dry N₂, unless otherwise noted. NMR (¹H and ¹³C) spectra were recorded on a Bruker AMX-400 MHz or Bruker Advance DRX-500 MHz and referenced to residual solvent unless otherwise noted. IR spectra were recorded with a Perkin-Elmer 1600 series-FTIR spectrometer and optical rotations were measured with a Jasco DIP-370 digital polarimeter using a 10-cm path length cell. Low-resolution mass spectral analysis were performed with a JOEL JMS-DX-303 HF mass spectrometer. Analytical TLC was performed on E. Merck silica gel 60 F254 plates and flash column chromatography was performed using the indicated solvents on E. Merck silica gel 60 (40-63 mm) or Sigma H-type silica gel (10-40 mm).

10

15

25

30

35

Procedure for the synthesis of 40 (as shown in Figure 17)

Trichloroacetimidate donor. The mixture of azidonitrates as shown in Figure 17 (1.66g, 4.41 mmol) was dissolved in CH₃CN (15 mL) and cooled to 0 °C. To the stirring solution was added Hunig's base (1.2 equiv., 0.925 mL) and benzene thiol (3.0 equiv., 1.35 mL). The reaction mixture was stirred at 0 °C for 1 hour and the ice bath was removed. After an additional 1 hour at room temperature, the reaction was concentrated under a stream of dry nitrogen. The resulting material was dissolved in a minimum amount of CHCl₃ and subjected to flash column chromatography (50% EtOAc/hexanes) to yield the hemiacetals (1.41, 97%). (Note 1: perform this flash in the hood, Note 2: isolate both anomers, which do separate on TLC/flash) The mixture of hemiacetals (1.41 g mg, 4.25 mmol) was dissolved in CH₂Cl₂ (8.5 mL) and trichloroacetonitrile (4.25 mL) was added, followed by K₂CO₃ (5.0 equiv., 2.93 g). The reaction stirred at room temperature overnight and was filtered through a plug of celite with additional methylene chloride. Concentration of the organic layer, followed by flash column chromatography (10->25% EtOAc/hexanes) gave the β-trichloroacetimidate (1.30 mg, 77%) as a yellow oil. (Note: α-anomer elutes first, then the β-anomer)

Pentenyl glycoside of α-Tn. The TCA-donor as shown in Figure 17 (1.30 g, 2.72 mmol) was dissolved in THF (0.2M, 13.6 mL) and pentenyl alcohol (5.0 equiv., 1.2 mL) and cooled to -10 °C (acetone-ice bath). A portion of TMSOTf (0.1 equiv., 0.049 mL) was added and the reaction stirred for 1 hour. Solid NaHCO₃ was added and the reaction was filtered through celite, concentrated and subjected to flash column chromatography (25% EtOAc/hexanes). (Note 1: The diastereomeric anomers do not separate. Their ratio is determined by ¹H NMR. Note 2: The starting materials and products co-elute by TLC - gradient TLC (10% first, then 50%) can be used to visualize the progress of the reaction.) The isolated glycosides were taken up in 10 mL of AcSH and stirred at rt for 2 days. Evaporation of the solvent by a stream of dry nitrogen followed by flash column chromatography (5% acetone/toluene -> 10% acetone/toluene) gave 620 mg of the α-glycoside (55%) and an undertermined amount of β-glycoside. (Note: Ethyl acetate/hexanes mixtures will also separate anomers, but actone/toluene was

determined to be superior.)

(

5

10

15

20

25

30

35

General procedure for olefination, 41. The preparation of enamide 41 (Globo-H) is representative of this procedure. The *n*-pentenyl glycoside 37 (58 mg, 0.0322 mmol) was dissolved in 10:10:1 MeOH:CH₂Cl₂:pyridine (3 mL, typically 0.05 M-0.01 M) and cooled to -78 °C. A stream of dry ozone was passed through the reaction mixture until a pale blue color persisted. The ozone source was removed and the reaction stirred at -78 °C for an additional 15 minutes, upon which time a stream of dry nitrogen was applied to remove excess ozone. Dimethyl sulfide (50 equivs., 0.118 mL) was added to the cooled mixture, the ice bath was removed and the reaction was allowed to stir at rt for 4 hours. The reaction was diluted with CH₂Cl₂ (10mL), washed with water (50mL), and back-extracted with additional CH₂Cl₂(2x10 mL). The combined organic layer was dried over anhydrous MgSO₄ and concentrated. The crude aldehyde was typically not purified, but was azeotroped dry with anhydrous benzene (3x3 mL) and used directly in the next step.

Phosphonate 33 (1.20 equivs., 14 mg) was dissolved in anhydrous THF (0.3 mL), cooled to -78 °C and tetramethyl guanidine (TMG) (1.25 equivs., 0.005 mL) was added dropwise. The reaction stirred at -78 for 30 minutes, followed by addition of the crude aldehyde (0.0322 mmol) in additional THF (2x0.3 mL, typically 0.1-0.01 M total reaction volume). The reaction was allowed to stir to rt overnight (10-15h), was extracted with EtOAc (10mL), washed with 0.05 M aqueous HCl (50mL) and back-extracted with additional EtOAc (2x10 mL). (Note: All the TMG must be removed prior to asymmetric hydrogenation.) The combined organic layer was dried over MgSO₄, concentrated and purified by flash column chromatography (75% EtOAc/hexanes -> 100 % EtOAc) to yield the desired enamide ester 41 as a single isomer. 72%, white foam; $R_f 0.85$ (100%) EtOAc); IR (CDCl₃ film) 3373, 2956, 2951, 1748, 1370, 1069 cm⁻¹; ¹H NMR (CDCl₃. 500 MHz) 8 6.65 (d, 1H, J=6.4 Hz), 6.44 (m, 1H), 6.07 (bs, 1H), 5.56 (d, 1H, J=3.1 Hz), 5.44 (d, 1H, J=3.4 Hz), 5.37 (d, 1H, J=3.3 Hz), 5.27 (dd, 1H, J=10.9, 3.0 Hz), 5.22 (d, 1H, J=2.6 Hz), 5.20-5.17 (m, 2H), 5.15 (d, 1H, J=2.1 Hz), 5.13 (d, 1H, J=4.9 Hz), 5.09 (dd. 1H, J=10.7, 7.3 Hz), 5.03 (dd, 1H, J=11.1, 3.3 Hz), 4.96 (dd, 1H, J=9.6, 3.5 Hz), 4.92 (dd, 1H, J=11.2, 3.4 Hz), 4.85 (dd, 1H, J=9.6, 8.0 Hz), 4.73 (dd, 1H, J=10.9, 2.5 Hz), 4.50-4.38 (m, 6H), 4.34 (t, 1H, J=6.2 Hz), 4.26-4.21 (m, 3H), 4.16-4.02 (m, 8H),

5 3.98 (d, 1H, J=2.0 Hz), 3.94 (t, 1H, J=6.4 Hz), 3.86-3.72 (m, 6H), 3.60-3.57 (m, 1H), 3.48-3.46 (m, 1H), 2.94-2.89 (m, 1H), 2.17-2.14 (m, 1H), 2.11-2.08 (m, 1H), 2.04 (s, 3H), 2.038 (s, 3H), 2.033 (s, 6H), 1.99 (s, 3H), 1.98 (s, 3H), 1.97 (s, 3H), 1.967 (s, 3H), 1.962 (s, 3H), 1.95 (s, 3H), 1.92 (s, 9H), 1.89 (s, 3H), 1.857 (s, 3H), 1.854 (s, 3H), 1.78 (s, 3H), 1.63-1.59 (m, 2H), 1.34 (s, 9H), 1.04 (d, 3H, J=6.5 Hz), 0.93-0.90 (m, 2H), -0.06 (s, 9H); ¹³C NMR (CDCl₃, 125 MHz) δ 172.20, 171.44, 170.94, 170.65, 170.52, 170.48, 170.44, 170.36, 170.29, 170.21, 170.17, 169.97, 169.63, 169.49, 169.31, 168.85, 164.78, 153.19, 134.70, 126.80, 102.35, 101.99, 101.26, 100.25, 99.12, 998.66, 94.21, 80.24, 76.88, 75.98, 73.61, 73.36, 73.08, 72.80, 72.56, 72.37, 71.81, 71.68, 71.46, 71.28, 70.78, 70.69, 70.67, 70.37, 70.06, 70.01, 68.92, 68.82, 67.99, 67.95, 67.54, 67.28, 66.94, 64.42, 62.14, 61.67, 61.29, 61.09, 60.92, 56.16, 28.12, 27.98, 24.52, 23.80, 23.03, 20.81, 20.73, 20.70, 20.68, 20.64, 20.60, 20.59, 20.54, 20.46, 20.40, 17.37, 17.24, 15.85, 15.48, 14.01, -1.58; HRMS (FAB) cald. for C₈₈H₁₂₈N₂O₅₁SiNa 2079.7145, found 2079.7174.

Lactose Enamide 34. 88%, white foam; R_f 0.45 (66% EtOAc/hexanes); IR (CDCl₃ film)
3407, 3146, 2954, 2898, 1752, 1654, 1233, 1167, 1055 cm⁻¹; ¹H NMR (CDCl₃, 400
MHz) δ 6.34 (m, 1H), 6.03 (bs, 1H), 5.22 (d, 1H, J=3.2 Hz), 5.07 (t, 1H, J=9.4 Hz), 4.99
(dd, 1H, J=10.3, 7.9 Hz), 4.83 (dd, 1H, J=10.5, 3.3 Hz), 4.77 (t, 1H, J=9.3 Hz), 4.37-4.33
(m, 3H), 4.20-4.11 (m, 3H), 4.08-4.00 (m, 3H), 3.82-3.65 (m, 5H), 3.49-3.46 (m, 1H),
3.39-3.34 (m, 1H), 2.16-2.14 (m, 1H), 2.11-2.09 (m, 1H), 2.04 (s, 3H), 2.02 (s, 3H), 1.94
(s, 3H), 1.93 (s, 3H), 1.92 (s, 6H), 1.84 (s, 3H), 1.62-1.59 (m, 2H), 1.34 (s, 9H), 0.94-0.89
(m, 2H), 0.05 (s, 9H); ¹³C NMR (CDCl₃, 100 MHz) δ 170.29, 170.21, 170.03, 169.94,
169.66, 169.50, 168.97, 164.77, 153.20, 134.70, 100.97, 100.31, 80.23, 76.17, 72.69,
72.51, 71.56, 70.87, 70.52, 68.95, 68.83, 66.47, 63.54, 61.88, 60.66, 33.81, 28.05, 27.92,
24.47, 20.73, 20.68, 20.57, 20.51, 20.39, 17.21, -1.60; HRMS (FAB) cald. for
C₄₃H₆₇NO₂₁SiNa 986.4013, found 986.4029.

Lewis enamide 43. 85%, white foam; R_f 0.45 (75% EtOAc/hexanes); IR (CDCl₃ film) 3371, 2965, 2956, 1746, 1371, 1231, 1069 cm⁻¹; ¹H NMR (CDCl₃, 500 MHz) δ 7.94 (d, 2H, J=7.2 Hz), 7.48 (t, 1H), 7.36 (t, 2H), 6.21 (t, 1H), 5.71 (m, 1H), 5.38 (d, 1H, J=3.2 Hz), 5.29 (dd, 1H, J=10.8, 8.8 Hz), 5.22-5.19 (m, 4H), 5.05-4.78 (m, 10H), 4.43 (dd, 1H,

J=14.1, 8.0 Hz), 4.36 (dd, 1H, J=9.7, 5.1 Hz), 4.27 (m, 1H), 4.15-4.30 (m, 5H), 3.86 (dd, 1H, J=10.1, 3.3 Hz), 3.78-3.71 (m, 3H), 3.62 (dd, 1H, J=9.8, 1.7 Hz), 3.30 (d, 1H, J=9.6 Hz), 3.18-3.08 (m, 1H), 3.04 (bm, 1H), 2.08 (s, 3H), 2.04 (s, 6H), 2.01 (s, 3H), 1.99 (s, 9H), 1.98 (s, 3H), 1.94 (s, 3H), 1.89 (s, 3H), 1.88 (s, 3H), 1.86 (s, 3H), 1.83 (s, 3H), 1.55-1.52 (m, 2H), 1.32 (s, 9H), 1.01-1.00 (m, 6H), 0.91-0.88 (m, 2H), -0.07 (s, 9H); ¹³C NMR (CDCl₃, 100 MHz) & 172.56, 172.40, 172.28, 172.11, 172.06, 172.04, 171.87, 171.77, 171.57, 171.43, 171.20, 171.06, 166.69, 166.32, 154.76, 136.48, 135.14, 131.29, 130.76, 130.19, 127.93, 102.82, 101.83, 101.72, 97.65, 97.11, 81.66, 75.40, 74.80, 74.55, 74.32, 74.01, 72.87, 72.65, 72.35, 72.28, 71.14, 70.66, 69.73, 69.28, 69.15, 69.02, 68.34, 66.43, 65.38, 64.92, 63.36, 62.05, 61.80, 59.87, 29.66, 29.62, 29.48, 25.83, 24.02, 22.51, 22.46, 22.31, 22.14, 22.12, 22.09, 22.003, 18.75, 17.30, 17.00, 15.63, -0.04; HRMS cald. for C₇₉H₁₁₂N₂O₄₁SiNa, found x.

Tn enamide 44. 75%, white foam; R_f 0.80 (100% EtOAc); IR (CDCl₃ film) 3340, 3071, 2954, 1715, 1663, 1498, 1369, 1218, 1162, 1049 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz) δ 6.38 (bs, 1H), 6.15 (bs, 1H), 5.26 (d, 1H, J=2.7 Hz), 5.02 (dd, 1H, J=11.5, 3.2 Hz), 4.79 (s, 1H), 4.48-4.42 (m, 1H), 4.16-4.12 (m, 2H), 4.03 (m, 1H), 3.99-3.92 (m, 2H), 3.62-3.56 (m, 1H), 3.40-3.35 (m, 1H), 2.26-2.16 (m, 2H), 2.03 (s, 3H), 1.92 (s, 3H), 1.86 (s, 3H), 1.84 (s, 3H), 1.67 (s, 3H), 1.33 (s, 9H), 0.94-0.9 (m, 2H), -0.06 (s, 9H); ¹³C NMR (CDCl₃, 100 MHz) δ 170.28, 170.11, 170.06, 164.70, 134.6, 108.73, 97.26, 80.31, 67.95, 67.05, 66.35, 63.51, 61.67, 47.46, 27.87, 27.49, 22.78, 20.43, 17.05, -1.74; HRMS (FAB) cald. for C₃₀H₅₁N₂O₁₃SiNa 675.3160, found 675.3124.

Fucosyl GM₁ enamide 42. 10-22%; R_f 0.25 (10% MeOH/EtOAc); ¹H NMR (MeOH, 500 MHz) 8 7.94 (d, 2H, J=7.2 Hz), 7.48 (t, 1H), 7.36 (t, 2H), 6.21 (t, 1H), 5.71 (m, 1H), 5.38 (d, 1H, J=3.2 Hz), 5.29 (dd, 1H, J=10.8, 8.8 Hz), 5.22-5.19 (m, 4H), 5.05-4.78 (m, 10H), 4.43 (dd, 1H, J=14.1, 8.0 Hz), 4.36 (dd, 1H, J=9.7, 5.1 Hz), 4.27 (m, 1H), 4.15-4.03 (m, 5H), 3.86 (dd, 1H, J=10.1, 3.3 Hz), 3.78-3.71 (m, 3H), 3.62 (dd, 1H, J=9.8, 1.7 Hz), 3.39-3.37 (m, 1H), 3.30 (bd, 1H, J=9.6 Hz), 3.18-3.08 (m, 1H), 3.04 (bm, 1H), 2.08 (s, 3H), 2.04 (s, 6H), 2.01 (s, 3H), 1.99 (s, 9H), 1.98 (s, 3H), 1.94 (s, 3H), 1.89 (s, 3H), 1.88 (s,

3H), 1.86 (s, 3H), 1.83 (s, 3H), 1.5-1.52 (m, 2H), 1.32 (s, 9H), 1.01-1.00 (m, 6H), 0.91-0.88 (m, 2H), -0.07 (s, 9H).

General procedure for asymmetric hydrogenation. Under an inert deoxygenated atmosphere, [(COD)Rh-((S, S)-Et-DuPHOS)]*OTf (0.005 mmol, 5 mol%) and the desired enamide ester (0.100 mmol) were dissolved in deoxygenated anhydrous THF (10 mL, 0.01 M) in a Fischer-Porter tube. The reaction vessel was pressurized with 50 psi of H₂ after three vacuum/H₂ cycles and stirred at 25 °C for 24-36 hours, or until the reaction turned from light orange to brown in color. The vessel was depressurized, the mixture concentrated and purified through a short plug of silica gel to yield the glycoamino acid.

15

20

25

30

35

10

Laetose glycoamino acid 35. 98%; R_f 0.45 (66% EtOAc/hexanes); 1 H NMR (C_6D_6 , 500 MHz) δ 5.54 (dd, 1H, J=10.4, 8.0 Hz), 5.48 (d, 1H, J=3.2 Hz), 5.39 (t, 1H, J=9.2 Hz), 5.21 (dd, 1H, J=6.2, 1.1 Hz), 5.12 (d, 1H, J=3.1 Hz), 5.09 (d, 1H, J=3.3 Hz), 4.54-4.51 (m, 2H), 4.33 (d, 1H, J=7.8 Hz), 4.19-4.06 (m, 6H), 3.74-3.58 (m, 2H), 3.49-3.40 (m, 1H), 3.38 (d, 1H, J=10.9 Hz), 3.23-3.16 (m, 2H), 1.96 (s, 3H), 1.94 (s, 3H), 1.90 (s, 3H), 1.74 (s, 3H), 1.73 (s, 3H), 1.64 (s, 3H), 1.51 (s, 3H), 1.45 (s, 9H), 0.91-0.88 (m, 2H), -0.10 (s, 9H); 13 C NMR (C_6D_6 , 100 MHz) δ 173.26, 170.44, 170.41, 170.18, 170.09, 169.35, 156.05, 102.98, 101.06, 79.73, 77.59, 74.13, 73.21, 72.73, 71.94, 71.19, 70.10, 69.58, 67.28, 63.76, 63.01, 61.25, 54.41, 34.76, 32.76, 28.62, 28.80, 25.75, 22.45, 21.18, 20.93, 20.84, 20.76, 20.57, 20.46, 20.15, 17.83, -1.29; HRMS (FAB) cald. for $C_{42}H_{67}$ NO₂₂SiNa 988.3870, found 988.3821.

Globo-H glycoamino acid 45. 98%; IR (CDCl₃ film) 3373, 2956, 2951, 1748, 1370, 1069 cm⁻¹; ¹H NMR (C₆D₆, 500 MHz) & 6.54 (d, 1H, J=6.5 Hz), 5.89 (d, 1H, J=3.5 Hz), 5.86 (d, 1H, J=3.1 Hz), 5.74-5.69 (m, 3H), 5.50-5.46 (m, 2H), 5.39-5.34 (m, 2H), 5.31 (dd, 1H, J=13.4, 0.7 Hz), 5.26 -5.19 (m, 2H), 5.15 (d, 1H, J=8.1 Hz), 5.09-5.06 (m, 3H), 4.82 (dd, 1H, J=10.9, 2.5 Hz), 4.74-4.72 (m, 1H), 4.70-4.69 (m, 1H), 4.66 (t, 1H), 4.58-4.49 (m, 3H), 4.45-4.41 (m, 2H), 4.37-4.33 (m, 2H), 4.24-4.22 (m, 2H), 4.20-4.12 (m, 3H), 4.04-4.01 (m, 2H), 3.91-3.86 (m, 3H), 3.71-3.69 (m, 2H), 3.64-3.57 (m, 2H), 4.43 (t, 1H), 3.28-3.27 (m, 1H), 3.23-3.21 (m, 1H), 2.38 (s, 3H), 2.30 (s, 3H), 2.19 (s, 3H), 2.01

5 (s, 3H), 1.98 (s, 3H), 1.93 (s, 3H), 1.92 (s, 3H), 1.90 (s, 3H), 1.89 (s, 3H), 1.88 (s, 3H), 1.84 (s, 3H), 1.82 (s, 3H), 1.75 (s, 3H), 1.73 (s, 3H), 1.71 (s, 3H), 1.68 (s, 3H), 1.64 (s, 3H), 1.61 (s, 3H), 1.46 (s, 3H), 1.45 (s, 9H), 1.26-1.22 (m, 2H), 1.08 (d, 3H, J=6.5 Hz), 0.89 (t, 2H), -0.10 (s, 9H); ¹³C NMR (CDCl₃, 125 MHz) & 172.87, 172.28, 171.52, 170.77, 170.61, 170.56, 170.45, 170.40, 170.29, 170.07, 169.75, 169.58, 169.44, 168.95, 155.38, 102.00, 101.29, 100.38, 99.21, 98.77, 94.31, 73.65, 73.34, 73.10, 72.61, 72.38, 71.84, 71.65, 71.58, 71.30, 70.81, 70.68, 70.17, 70.06, 69.59, 69.09, 68.85, 68.05, 67.99, 67.56, 67.30, 64.46, 63.64, 62.16, 61.74, 61.35, 61.12, 60.96, 56.11, 53.45, 32.30, 29.65, 28.96, 28.29, 28.10, 23.11, 21.69, 20.88, 20.85, 20.80, 20.76, 20.72, 20.67, 20.62, 20.55, 20.48, 17.31, 15.88, -1.55; HRMS (FAB) cald. for C₈₈H₁₃₀N₂O₅₁SiNa 2081.7302, found 2081.7247.

Lewis glycoamino acid 42. 99%; IR (CDCl₃ film) cm⁻¹; ¹H NMR (CDCl₃, 400 MHz) δ 8.24 (d, 2H, J=8.4 Hz), 7.27-7.18 (m, 3H), 5.83 (dd, 1H, J=10.1, 8.0), 5.78 (d, 1H, J=3.2 Hz), 5.76 (d, 1H, J=3.0 Hz), 5.74-5.71 (m, 2H), 5.69 (d, 1H, J=3.2 Hz), 5.66 (d, 1H, J=3.3Hz), 5.55 (d, 1H, J=3.3 Hz), 5.50 (m, 1H), 5.47 (d, 1H, J=3.8 Hz), 5.41-5.27 20 (m, 5H), 4.93 (d, 1H, J=7.8 Hz), 4.86 (d, 1H, J=8.6 Hz), 4.77-4.70 (m, 2H), 4.62 (d, 1H, J=7.8 Hz), 4.56-4.53 (m, 1H), 4.47-4.35 (m, 5H), 4.34-4.25 (m, 4H), 4.11-4.00 (m, 5H), 3.92-3.89 (m, 1H), 3.75-3.65 (m, 3H), 3.64-3.61 (t, 1H), 3.28-2.24 (m, 1H), 2.22 (s, 3H), 2.21 (s, 3H), 1.93 (s, 3H), 1.90 (s, 3H), 1.82 (s, 3H), 1.78 (s, 3H), 1.73 (s, 3H), 1.72 (s, 3H), 1.66 (s, 3H), 1.65 (s, 3H), 1.64 (s, 3H), 1.63 (s, 3H), 1.51 (s, 3H), 1.50 (d, 3H, J=6.5 Hz), 1.46 (s, 9H), 1.38 (d, 3H, J=6.5 Hz), 1.33-1.27 (m, 2H), 1.22-1.21 (m, 2H), 0.91- $0.80 \text{ (m, 4H)}, -0.12 \text{ (s, 9H)}; ^{13}\text{C NMR (C}_6\text{D}_6, 100 \text{ MHz)} \delta 172.86, 171.05, 170.59,$ 170.52, 170.46, 170.37, 170.26, 170.01, 169.97, 169.90, 165.44, 155.61, 133.24, 130.40, 101.59, 100.98, 100.71, 97.09, 96.21, 79.22, 76.42, 74.77, 74.04, 73.69, 72.03, 71.65, 71.14, 70.14, 69.20, 68.66, 68.55, 68.20, 67.99, 67.25, 65.67, 64.56, 63.23, 62.41, 61.33, 60.94, 58.23, 53.84, 39.12, 32.25, 29.30, 28.41, 22.88, 22.01, 20.93, 20.66, 20.58, 20.47, 20.40, 20.18, 20.02, 17.40, 16.36, 15.94, -1.62; HRMS (FAB) cald. for C₇₉H₁₁₄N₂O₄₁SiNa 1797.6558, found 1797.6528.

Tn glycoamino acid 43. 99%; IR (CDCl₃ film) 3362, 2954, 2990, 2871, 1749, 1716, 1683, 1668, 1520, 1369, 1249, 1164, 1047 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz) δ 5.66 (d, 1H, J=9.3 Hz), 5.24 (d, 1H, J=2.8 Hz), 5.03 (dd, 1H, J=11.4, 3.3 Hz), 4.98 (d, 1H, J=8.1), 4.73 (d, 1H, J=3.3 Hz), 4.48-4.42 (m, 1H), 4.17-4.07 (m, 3H), 4.05-3.93 (m, 3H), 3.59-3.54 (m, 1H), 3.33-3.27 (m, 1H), 2.04 (s, 1H), 1.93 (s, 3H), 1.87 (s, 3H), 1.86 (s, 3H), 1.73-1.70 (m, 1H), 1.57-.150 (m, 3H), 1.31 (s, 9H), 0.91-0.87 (m, 2H), -0.06 (s, 9H); ¹³C NMR (C₆D₆, 100 MHz) δ 173.44, 171.04, 170.57, 170.35, 169.89, 156.06, 98.59, 79.82, 69.29, 68.30, 68.07, 67.43, 64.01, 62.33, 54.33, 48.63, 32.87, 29.05, 28.76, 23.24, 22.82, 20.89, 20.66, 20.47, 17.85, -1.28; HRMS (FAB) cald. for C₃₀H₅₃N₂O₁₃SiNa 677.3316, found 677.3352.

15

20

35

General procedure for N-Boc deprotection. The desired glycoamino acid (0.100 mmol) was dissolved in CH₂Cl₂ (3.0mL) with stirring. Trifluoroacetic acid (TFA) (3.0mL) was added dropwise and the reaction stirred at rt for 1 hour. The mixture was then concentrated with a stream of dry N₂ and azeotroped with anhydrous benzene (2x5mL) to give the crude amine as its TFA salt which was typically used without further purification.

General procedure for TSE ester deprotection. The desired glycoamino acid (0.100 mmol) was dissolved in THF (1.0-3.0 mL) and cooled to 0 °C. A 1.0 M solution of TBAF in THF (0.250 mmol, 2.5 equivs.) was added dropwise, the ice bath removed and the reaction stirred at rt for 1-2 hour, as judged by TLC. (Note: prolonged reaction times, i.e. >10 h, may result in deacetylation.) The reaction mixture was diluted with CH₂Cl₂ (10mL), washed with 0.05M aqueous HCL (50 mL), and back-extracted with additional CH₂Cl₂ (2x10 mL). The combined organic layer was dried over anhydrous Mg₂SO₄ and concentrated. The crude acid was typically used without further purification. Acid 36: ¹H NMR (CDCl₃, 400 MHz) 8 5.22 (d, 1H, J=2.8 Hz), 5.07 (t, 1H, J=9.3 Hz), 4.98 (dd, 1H, J=10.4, 5.9 Hz), 4.84 (dd, 1H, J=10.4, 3.5 Hz), 4.75 (dd, 1H, J=9.5, 8.0 Hz), 4.42-4.35 (m, 2H), 4.34-4.31 (m, 1H), 4.15-4.14 (m, 1H), 4.03-3.94 (m, 4H), 3.77-3.65 (m, 5H), 3.49-3.45 (m, 1H), 3.37-3.33 (m, 1H), 3.10-3.07 (m, 1H), 2.03 (s, 3H), 2.00 (s, 3H), 1.96 (s, 3H), 1.94 (s, 3H), 1.92 (s, 3H), 1.84 (s, 3H), 1.51-1.46 (m, 4H), 1.32 (s, 9H).

(

General procedure for BOP reagent promoted peptide coupling. The desired amine and acid (equimolar amounts) were azeotroped together with anhydrous benzene and dried under high vacuum. The mixture was dissolved in CH₂Cl₂ (0.1-0.05M), BOP reagent (1.25 equivs) was added and the solution cooled to 0 °C over 15 minutes. A dropwise addition of Hunig's base (15 equivs.) was followed by removal of the ice bath. The reaction stirred at rt for 2-4 h, as judged by TLC. Concentration of the reaction mixture was followed by purification by flash column chromatography. In cases where bi-product HMPA was difficult to remove, the peptide was subjected to sephadex purification (LH-20, MeOH).

15

20

25

30

35

N-Boc Tn with mercatoacetamide spacer. 54%, colorless oil; R_f 0.35 (10% MeOH/EtOAc); IR (CDCl₃ film) 3303, 3078, 2974, 2935, 2872, 1748, 1703, 1692, 1658, 1535, 1440, 1369, 1245, 1166 cm⁻¹; ¹H NMR (MeOH, 400 MHz) δ 5.40 (d, 1H, J=2.9 Hz), 5.13 (dd, 1H, J=11.6, 3.2 Hz), 4.42 (dd, 1H, J=11.5, 3.5 Hz), 4.23 (t, 1H, J=6.7 Hz), 4.14-4.05 (m, 2H), 3.96 (m, 1H), 3.74-3.69 (m, 1H), 3.60 (s, 2H), 3.49-4.44 (m, 1H), 3.26-3.13 (m, 5H), 2.36 (s, 3H), 2.12 (s, 3H), 2.02 (s, 3H), 1.94 (s, 3H), 1.93 (s, 3H), 1.67-1.59 (m, 6H), 1.43 (s, 9H); ¹³C NMR (CDCl₃, 100 MHz) δ 195.59, 172.53, 170.78, 170.54, 170.39, 170.30, 168.51, 155.62, 97.59, 79.95, 68.29, 67.96, 67.26, 66.38, 61.83, 60.28, 54.58, 47.62, 36.47, 35.97, 33.02, 31.92, 30.18, 29.14, 28.37, 28.17, 23.07, 22.35, 20.68, 20.64; HRMS (MALDI) cald. for C₃₂H₅₂N₄O₁₄SNa 771.3093, found 771.3070.

Le^Y/Tn dipeptide 51. 86%, white film; R_f 0.65 (20% MeOH/EtOAc); ¹H NMR (MeOH, 400 MHz) δ 8.06 (d, 2H, J=7.4 Hz), 7.63 (t, 1H), 7.51 (t, 2H), 5.55 (d, 1H, J=3.3 Hz), 5.40 (d, 1H, 2.7 Hz), 5.38 (d, 1H, J=2.7 Hz), 5.32 (d, 1H, J=3.3 Hz), 5.29 (d, 1H, J=4.1 Hz), 5.26 (d, 1H), 5.21-5.12 (m, 5H), 5.01 (q, 1H), 4.93 (m, 2H), 4.90 (m, 2H), 4.79 (d, 1H, J=10.8 Hz), 4.71 (d, 1H), 4.64 (d, 1H), 4.50-4.01 (m, 15H), 3.88-3.60 (m, 7H), 3.60 (s, 2H), 3.51-3.42 (m, 2H), 3.20-3.13 (m, 5H), 2.36 (s, 3H), 2.20 (s, 3H), 2.16 (s, 3H), 2.15 (s, 3H), 2.13 (s, 3H), 2.11 (s, 3H), 2.09 (s, 3H), 2.08 (s, 3H), 2.07 (s, 3H), 2.05 (s, 3H), 2.02 (s, 6H), 1.97 (s, 6H), 1.95 (s, 3H), 1.94 (s, 9H), 1.91 (s, 3H), 1.68-1.63 (m, 8H), 1.44 (s, 9H), 1.15 (d, 3H, J=6.3 Hz), 1.14 (d, 3H, J=6.3 Hz); ¹³C NMR (MeOH, 100

MHz) δ 196.26, 175.39, 174.30, 173.75, 173.22, 172.80, 172.72, 172.56, 172.51, 172.44, 172.39, 172.30, 172.19, 172.12, 171.85, 171.75, 171.64, 170.76, 166.77, 158.24, 134.92, 131.29, 131.10, 129.99, 103.53, 102.80, 101.77, 99.12, 97.81, 97.30, 80.90, 78.50, 75.88, 75.11, 74.65, 74.32, 73.01, 72.70, 72.56, 72.47, 72.18, 71.91, 70.86, 69.84, 69.62, 69.42, 69.09, 68.90, 67.84, 66.43, 65.43, 63.50, 63.21, 62.57, 61.68, 56.42, 55.96, 54.89, 38.21, 37.74, 37.12, 33.89, 32.72, 30.42, 30.21, 30.01, 28.92, 23.72, 23.44, 22.93, 22.73, 21.32, 21.13, 20.97, 20.86, 20.74, 20.60, 16.67, 16.29; HRMS (MALDI) cald. for C₁₀₁H₁₄₄N₆O₅₂SiNa 2327.8421, found 2327.8536.

N-Boc Globo-H/Le^y/Tn tripeptide. 64%, white film; R_f 0.45 (10% MeOH/EtOAc); ¹H

NMR (MeOH, 400 MHz) δ 8.05 (d, 2H, J=7.4 Hz), 7.63 (t, 1H), 7.50 (t, 2H), 5.64 (d, 1H, J=2.8 Hz), 5.55 (d, 1H, J=3.6 Hz), 5.43 (d, 1H, J=3.2 Hz), 5.40 (d, 1H, J=2.4 Hz), 5.37 (d, 1H, J=2.5 Hz), 5.32-5.25 (m, 7H), 5.23-5.12 (m, 10H), 5.08-5.05 (m, 2H), 5.00 (d, 1H, J=7.5 Hz), 4.96 (d, 1H, J=3.1 Hz), 4.94 (m, 2H), 4.90 (m, 2H), 4.86 (m, 2H), 4.82-4.77 (m, 3H), 4.72-4.70 (m, 3H), 4.64-4.59 (m, 3H), 4.51-4.35 (m, 9H), 4.32-3.92 (m, 31H), 3.86-3.67 (m, 13H), 3.60 (s, 2H), 3.59 (m, 1H), 3.51 (m, 1H), 3.47-3.44 (m, 2H), 3.24-3.18 (m, 5H), 2.36 (s, 3H), 2.19 (s, 3H), 2.17 (s, 3H), 2.16 (s, 3H), 2.15 (s, 3H), 2.14 (s, 3H), 2.138 (s, 3H), 2.133 (s, 3H), 2.12 (s, 3H), 2.116 (s, 3H), 2.115 (s, 3H), 2.10 (s, 3H), 2.096 (s, 3H), 2.090 (s, 3H), 2.08 (s, 3H), 2.07 (s, 3H), 2.05 (s, 6H), 2.04 (s, 3H), 2.039 (s, 3H), 2.031 (s, 9H), 2.02 (s, 6H), 2.00 (s, 3H), 1.97 (s, 3H), 1.96 (s, 3H), 1.94 (s, 3H), 1.93 (s, 3H), 1.91 (s, 3H), 1.86 (s, 3H), 1.67-1.52 (m, 16H), 1.43 (s, 9H), 1.32 (d, 3H), 1.16-1.13 (m, 6H).

N-Ac capped Globo-H/Le^y/Tn tripeptide 53. 95%, white film; R_f 0.35 (10% MeOH/EtOAc); ¹H NMR (CDCl₃, 500 MHz) δ 8.00 (d, 2H, J=7.3 Hz), 7.59 (t, 1H), 7.47 (t, 2H), 6.70 (d, 1H, J=6.3 Hz), 6.61 (d, 1H, J=9.8 Hz), 5.56 (d, 1H, J=2.9 Hz), 5.47 (d, 1H, J=2.1 Hz), 5.44 (d, 1H, J=3.4 Hz), 5.37 (d, 1H, J=3.2 Hz), 5.34 (d, 1H, J=2.7 Hz), 5.32-5.28 (m, 7H), 5.28 (d, 1H), 5.22-5.19 (m, 3H), 5.15-5.10 (m, 6H), 5.08 (m, 1H), 5.06 (m, 1H), 5.01-4.87 (m, 13H), 4.85-4.82 (m, 1H), 4.55-4.35 (m, 15H), 4.26-4.23 (m, 4H), 4.15-4.02 (m, 20H), 3.98-3.93 (m, 5H), 3.88-3.72 (m, 12H), 3.66 (m, 1H), 3.59-3.58 (m, 1H), 3.52 (s, 2H), 3.49-3.38 (m, 5H), 3.20 (m, 6H), 3.02-2.98 (m, 1H), 2.37 (s, 3H),

2.17 (s, 3H), 2.139 (s, 6H), 2.133 (s, 6H), 2.12 (s, 6H), 2.11 (s, 3H), 2.10 (s, 3H), 2.08 (s, 12H), 2.07 (s, 3H), 2.06 (s, 12H), 2.05 (s, 3H), 2.046 (s, 3H), 2.041 (s, 3H), 2.03 (s, 3H), 2.019 (s, 3H), 2.015 (s, 9H), 1.98 (s, 3H), 1.97 (s, 6H), 1.955 (s, 3H), 1.951 (s, 3H), 1.94 (s, 3H), 1.92 (s, 3H), 1.88 (s, 3H), 1.76-1.57 (m, 10H), 1.44-1.41 (m, 4H), 1.38-1.30 (m, 4H), 1.23-1.20 (m, 2H), 1.13-1.09 (m, 9H).

10

(

Fully Deprotected Globo-H/Le^Y/Tn tripeptide 54. 98%, white film; ¹H NMR (D₂O, 500 MHz) δ 5.30 (s, 1H, J=2.8 Hz), 5.25 (d, 1H, J=3.7 Hz), 5.13 (d, 1H, J=3.5 Hz), 4.91-4.87 (m, 3H), 4.75-4.74 (m, 1H), 4.63 (d, 1H, J=7.4 Hz), 4.57-4.48 (m, 3H), 4.41-4.38 (m, 2H), 4.31 (m, 1H), 4.29-4.24 (m, 6H), 4.18-4.12 (m, 3H), 4.05-3.56 (m, 58H), 3.50-3.46 (m, 3H), 3.32-3.24 (m, 5H), 2.06 (s, 3H), 2.05 (s, 3H), 2.04 (s, 6H), 1.82-1.65 (m, 14H), 1.47-1.42 (m, 6H), 1.29 (d, 3H, J=6.5 Hz), 1.25 (d, 3H, J=7.4 Hz), 1.23 (d, 3H, J=7.0 Hz); ¹³C NMR (D₂O, 125 MHz) δ 175.11, 174.85, 174.81, 174.68, 174.44, 174.25, 174.13, 171.64, 164.59, 104.37, 103.71, 103.25, 102.78, 102.39, 102.44, 100.83, 100.59, 99.80, 99.67, 98.96, 97.34, 82.76, 79.18, 78.65, 77.55, 76.74, 76.50, 75.88, 75.78, 75.45, 75.24, 75.18, 75.00, 74.90, 73.96, 73.47, 73.35, 72.50, 72.32, 72.24, 72.09, 71.31, 71.24, 70.53, 70.44, 70.40, 70.16, 70.11, 69.90, 69.57, 69.50, 69.12, 68.93, 68.87, 68.66, 68.42, 68.22, 68.12, 68.08, 67.29, 67.17, 61.85, 61.68, 61.37, 61.34, 61.20, 60.74, 60.48, 60.19, 56.57, 54.02, 52.02, 50.41, 41.31, 37.09, 31.07, 30.79, 28.69, 28.64, 28.36, 22.72, 22.64, 22.40, 22.36, 22.25, 22.05, 22.01, 21.97, 21.93, 21.87, 21.84, 15.84, 15.70.

25

3) Preparation of Polycarbohydrate (globo H, Le^y, Tn) cluster-KLH conjugate using bifunctional cross linker method:

Polycarbohydrate (globo H, Le^y, Tn) cluster is conjugated as described below using maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) which is a heterobifunctional reagant. At neutral pH it crosslinks amino groups with succinimide and then with thiol groups with maleimide. The thiol group is provided by the cysteine residue of peptide backbone of cluster and the amino groups by the N-terminal and lysine side-chain of the KLH. After linkage MBS to KLH the unreacted MBS is purified by column and crosslinked to cysteine on the synthetic polycarbohydrate cluster. The unbound antigen is removed by passage of the reaction mixture through a CentriPrep 30 filter with a 30,000

molecular weight cut-off. The epitope ratio is then calculated by estimation of protein content by standard method and carbohydrate by high-pH anion-exchange chromatography with pulsed amperometric detection (HPAEC-PAD) method.

D. Example 4: Immunological Studies

It will be appreciated that the inventive glycoconjugates and glycopeptides, as provided herein, are useful for the treatment of cancer and are useful for inducing an antibody response in a subject. Typical protocols for the use of such glycoconjugates and glycopeptides are described in more detail below, and are also detailed in certain references incorporated herein.

15

20

25

10

1) Immunization of Mice

Groups of mice (CB6F1 female; 6 weeks of age) obtained from Jackson Laboratory, Bar Harbor, ME, are immunized subcutaneously with Polycarbohydrate cluster-KLH (globo H, Le^y, Tn) containing equivalent to 3 µg of total carbohydrate only (the quantity of KLH varied depending on the epitope density) mixed with 10 µg of immunological adjuvant QS-21, a saponin derivative from the bark of the *Quillaja saponaria* Molina tree (Ragupathi *et al. Angew. Chem. Int. Ed. Engl.* 1997, 36, 125) (Aquila, Worcester, MA) at 0, 1 and 2 weeks and bled 10 days after the 3rd immunization. The presence of antibody is assayed by an enzyme linked immunosorbent assay (ELISA) as described previously (Ragupathi *et al. Angew. Chem. Int. Ed. Engl.* 1997, 36, 125) using the appropriate target antigen (e.g., globo H-ceramide, Le^y ceramide and/or Tn(c)-pamcys as target antigen). The cell surface reactivity can then be tested, for example, the cell surface reactivity of anti- globo H, Le^y, and Tn antibodies are tested on globo H, Le^y, Tn positive cell-lines by flow cytometry assays.

30

35

1

2) Serological Analysis:

ELISA: Enzyme-linked immunosorbent assays (ELISAs) are performed as described (Ragupathi G, Park TK, Zhang S, Kim I-J, Graber L, Adluri S, Lloyd KO, Danishefsky SJ, Livingston PO. Immunization of mice with a fully synthetic globo H antigens results in antibodies against human cancer cells; A combined chemical-

ĺ,

15

20

immunological approach to the fashioning of an anticancer vaccine. Angew. Chem. Int. Ed. Engl. 1997, 36, 125). Serially diluted antiserum is added to wells coated with antigen (0.1 μg) and incubated for 1 h at room temperature. Goat anti-mouse IgM or IgG conjugated with alkaline phosphatase serve as secondary antibodies. Absorbance is measured at 414 nm. The antibody titer is defined as the highest serum dilution showing an absorbance 0.1 or greater above that of normal mouse sera.

3) Flow Cytometry:

Appropriate cells (e.g., cells from the globo H and Le^y-positive breast cancer cell line MCF-7 and Colon cancer cell-line LS-C) are used as target. Single cell suspensions of 2 X 10⁵ cells/tube are washed in PBS with 3% fetal calf serum and 0.01 M NaN3 and incubated with 20 µl of 1:20 diluted antisera or mAb VK-9 for 30 min on ice. After washing the cells twice with 3% FCS in PBS, 20 µl of 1:15 goat anti-mouse IgM or IgG-labeled with fluorescein-isothiocyanate (FITC) is added, mixed and incubated for 30 min. After wash, the positive population and mean fluorescence intensity of stained cells are analyzed by flow cytometry (EPICS-Profile II).

ĺ

Claims ·

1. A compound having the structure:

10

15

wherein R is hydrogen; substituted or unsubstituted alkyl; alkenyl; aryl;
-CH₂CH(CO₂R')(NHR"), wherein R' or R" are each indpendently hydrogen, protecting group, substituted or unsubstituted alkyl, a linker, aryl, peptide, protein or lipid; or NHR", wherein R" is a protein, peptide, or lipid linked to N directly or through a crosslinker; wherein n is 0-8; and wherein A is a carbohydrate domain having the structure:

20

25

ĺ

wherein a, b, c, d, e, f, g, h, i, x, y and z are independently 0, 1, 2 or 3, with the proviso that x, y and z are not simultaneously 0; wherein R₀ is hydrogen, a linear or branched chain alkyl, acyl, arylalkyl or aryl group; wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈ and R₉ are each independently hydrogen, OH, ORⁱ, NH₂, NHCORⁱ, F, CH₂OH, CH₂ORⁱ, a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein Rⁱ is

hydrogen, CHO, COORⁱⁱ, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group or a saccharide moiety having the structure:

- wherein Y and Z are independently NH or O; wherein k, l, r, s, t, u, v and w are each independently 0, 1 or 2; wherein R₁₀, R₁₁, R₁₂, R₁₃, R₁₄ and R₁₅ are each independently hydrogen, OH, ORⁱⁱⁱ, NH₂, NHCORⁱⁱⁱ, F, CH₂OH, CH₂ORⁱⁱⁱ, or a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein R₁₆ is hydrogen, COOH, COORⁱⁱ, CONHRⁱⁱ, a substituted or unsubstituted linear or branched chain alkyl or aryl group; wherein Rⁱⁱⁱ is hydrogen, CHO, COOR^{iv}, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group; and wherein Rⁱⁱ and R^{iv} are each independently H, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group, with the proviso that if A is KH-1, N3, Globo-H,
 glycophorin, Tn, TF, STN, (2,3)ST, 2,6-STn or Le^y, and A is α-O-linked, then n is at least 1.
 - 2. The compound of claim 1, wherein R is allyl.
- 25 3. The compound of claim 1, wherein n is 1 and R is allyl.
 - 4. The compound of claim 1, wherein n is 2, and R is allyl.
- 5. The compound of claim 1, wherein R is NHR", and wherein the protein R" is KLH or Bovine Serine Albumin, whereby said compound is a glycoconjugate.

ţ

- The compound of claim 1, wherein R is NHR", and wherein the lipid R" is 6. PamCys, whereby said compound is a glycoconjugate.
- The compound of claim 1, wherein R is CH₂CH(CO₂R')(NHR") and the resulting 7. glycopeptide has the structure: 10

The compound of claim 7 wherein n is 3. 8:

15

The compound of claim 7, wherein R' and R" are each hydrogen or a protecting 9. group.

The compound of claim 9, wherein R' and R" are each protecting groups 10. independently selected from the group consisting of Fmoc, acetyl, Boc, t-butyl and TSE. 20

The compound of claim 1, 4, 7 or 8, or the glycoconjugate of claim 5 or 6, 11. wherein the carbohydrate determinant is selected from the group consisting of Globo-H, fucosyl GM1, KH-1, glycophorin, N3,Tn, TF, STN, (2,3)ST, 2,6-STn, and Ley.

25

12. The compound or glycoconjugate of claim 11, wherein A is the carbohydrate determinant fucosyl GM1 having the structure:

(:


wherein each occurrence of R' is independently hydrogen or a protecting group; and wherein each occurrence of R" is independently hydrogen or a nitrogen protecting group.

10 13. The compound or glycoconjugate of claim 11, wherein A is the carbohydrate determinant Globo-H having the structure:

- wherein each occurrence of R' is independently hydrogen or a protecting group, and wherein R" is hydrogen or a nitrogen protecting group.
 - 14. A method for the synthesis of complex carbohydrates comprising:
 - (a) providing a carbohydrate acceptor having a reducing end alkenyl group;
- · 20 (b) providing a suitable donor compound; and

- 5 (c) coupling said donor and acceptor under conditions to generate an alkenyl glycoside.
 - 15. The method of claim 14, wherein the step of providing a carbohydrate acceptor having a reducing end alkenyl group comprises providing an acceptor having the structure:

wherein P is a protecting group and n is 0-8, and wherein the step of providing a suitable donor compound comprising providing a donor having the structure:



15

20

10

wherein n is 0-8, and wherein P is a protecting group.

16. The method of claim 14, wherein the step of providing a carbohydrate acceptor having a reducing end alkenyl group comprises providing an acceptor having the structure:

wherein P is a protecting group and n is 0-8, and wherein the step of providing a suitable donor compound comprising providing a donor having the structure:

10

20

5

wherein n is 0-8 and P is a protecting group.

17. A method for the synthesis of a glycoamino acid comprising the steps of:

(a) providing an alkenyl glycoside having the structure:

and reacting said alkenyl glycoside under suitable conditions to generate an enamide ester having the structure:

(b) reacting said enamide ester under suitable conditions to generate a glycoamino acid having the structure:

wherein, for each of the structures above, n is 0-8, wherein A is a carbohydrate domain having the structure:

$$R_0$$
 R_0
 R_0

15

5

10

wherein a, b, c, d, e, f, g, h, i, x, y and z are independently 0, 1, 2 or 3, with the proviso that x, y and z are not simultaneously 0; wherein R₀ is hydrogen, a linear or branched chain alkyl, acyl, arylalkyl or aryl group; wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈ and R₉ are each independently hydrogen, OH, OR¹, NH₂, NHCOR¹, F, CH₂OH, CH₂OR¹, a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein R¹ is hydrogen, CHO, COOR¹¹, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group or a saccharide moiety having the structure:

25

$$R_0$$
 Z
 R_{12}
 R_{12}
 R_{15}
 R_{16}
 R_{16}
 R_{16}

10

15

Ĺ

wherein Y and Z are independently NH or O; wherein k, l, r, s, t, u, v and w are each independently 0, 1 or 2; wherein R₁₀, R₁₁, R₁₂, R₁₃, R₁₄ and R₁₅ are each independently hydrogen, OH, ORⁱⁱⁱ, NH₂, NHCORⁱⁱⁱ, F, CH₂OH, CH₂ORⁱⁱⁱ, or a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein R₁₆ is hydrogen, COOH, COORⁱⁱ, CONHRⁱⁱ, a substituted or unsubstituted linear or branched chain alkyl or aryl group; wherein Rⁱⁱⁱ is hydrogen, CHO, COOR^{iv}, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group; and wherein Rⁱⁱ and R^{iv} are each independently H, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group;


and wherein for the glycoamino acid structure R' and R" are each independently protecting group or hydrogen.

- 18. The method of claim 17, wherein the carbohydrate determinant is selected from the group consisting of Globo-H, fucosyl GM1, KH-1, glycophorin, STN, (2,3)ST, Le^y, N3, Tn, 2,6-STn, and TF.
- 19. The method of claim 18, wherein A is a carbohydrate determinant having the structure:

30

wherein each occurrence of R' is independently hydrogen or a protecting group; and wherein each occurrence of R" is independently hydrogen or a nitrogen protecting group.

10 20. The method of claim 18, wherein A is a carbohydrate determinant having the structure:



- wherein each occurrence of R' is independently hydrogen or a protecting group, and wherein R" is hydrogen or a nitrogen protecting group.
 - 21. The method of claim 17, wherein the step of reacting an alkenyl glycoside under suitable conditions to generate an enamide ester comprises reacting an alkenyl glycoside

25

30

- first under oxidative cleavage conditions and second under olefination conditions in the presence of base and phosphonate to generate an enamide ester.
 - 22. The method of claim 21, wherein said oxidative cleavage conditions comprise ozonolysis, and wherein the base is tetramethylguanidine.
 - 23. The method of claim 21, wherein said oxidative cleavage conditions are OsO₄ and periodate, or OsO₄ and Pb(OAc)₄, and wherein the base is lithium t-butoxide or lithium hexamethyl disilylazide.
- 15 24. The method of claim 17, wherein the step of reacting said enamide ester under suitable conditions to generate a glycoamino acid comprises reacting said enamide ester under hydrogenation conditions and subsequent reaction under deprotection conditions to generate a glycoamino acid.
- 20 25. The method of claim 24, wherein said hydrogenation is achieved via asymmetric hydrogenation.
 - 26. The method of claim 25, wherein said asymmetric hydrogenation is achieved by utilizing an ethyl DuPHOS catalyst precursor.
 - 27. A multi-antigenic glycopeptide comprising a peptidic backbone made up of at least three amino acids, wherein one or more of said amino acids are substituted with an n-alkyl glycosidic moiety having the structure:

wherein each occurrence of A is a carbohydrate determinant having the structure:

$$\begin{bmatrix} R_7 & & & \\ & & &$$

wherein a, b, c, d, e, f, g, h, i, x, y and z are independently 0, 1, 2 or 3, with the proviso that x, y and z are not simultaneously 0; wherein R₀ is hydrogen, a linear or branched chain alkyl, acyl, arylalkyl or aryl group; wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈ and R₉ are each independently hydrogen, OH, OR¹, NH₂, NHCOR¹, F, CH₂OH, CH₂OR¹, a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein R¹ is hydrogen, CHO, COOR¹¹, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group or a saccharide moiety having the structure:

15

20

5

10

wherein Y and Z are independently NH or O; wherein k, l, r, s, t, u, v and w are each independently 0, 1 or 2; wherein R₁₀, R₁₁, R₁₂, R₁₃, R₁₄ and R₁₅ are each independently hydrogen, OH, ORⁱⁱⁱ, NH₂, NHCORⁱⁱⁱ, F, CH₂OH, CH₂ORⁱⁱⁱ, or a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein R₁₆ is hydrogen, COOH, COORⁱⁱ, CONHRⁱⁱ, a substituted or unsubstituted linear or branched

chain alkyl or aryl group; wherein Rⁱⁱⁱ is hydrogen, CHO, COOR^{iv}, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group; and wherein Rⁱⁱ and R^{iv} are each independently H, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group;

wherein each occurrence of n is independently 0-8, whereby, if for each occurrence of n, n = 0, at least one occurrence of A has a different structure from other occurrences of A; and wherein the n-alkyl glycosidic moiety is either α - or β -linked to an amino acid.

28. The glycopeptide of claim 27, wherein said glycopeptide is a construct having the structure:

wherein the linker is either a free carboxylic acid, (carboxamido)alkyl carboxamide, MBS, primary carboxamide, mono- or dialkyl carboxamide, diarylcarboxamide, linear or branched chain (carboxy)alkyl carboxamide, linear or branched chain (alkoxycarbonyl)alkyl-carboxamide, linear or branched chain (carboxy)arylalkylcarboxamide, branched chain linear or (alkoxycarbonyl)alkylcarboxamide, an oligoester fragment comprising from 2 to about 20 hydroxy acyl residues, a peptidic fragment comprising from 2 to about 20 amino acyl residues, or a linear or branched chain alkyl or aryl carboxylic ester; wherein the carrier is a protein or lipid; wherein m is 1, 2 or 3; wherein RA, RB and RC are each independently H or methyl; and wherein RD, RE and RF are each independently an alkyl glycosidic moiety having the structure:

10

15

wherein each occurrence of A is independently selected from a carbohydrate domain having the structure:

$$R_0$$
 R_0 R_0

.10

15

5

wherein a, b, c, d, e, f, g, h, i, x, y and z are independently 0, 1, 2 or 3, with the proviso that x, y and z are not simultaneously 0; wherein the carbohydrate domain is linked to the respective amino acyl or hydroxy acyl residue by substitution of a side group substituent selected from the group consisting of OH, COOH and NH₂; wherein R₀ is hydrogen, a linear or branched chain alkyl, acyl, arylalkyl or aryl group; wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈ and R₉ are each independently hydrogen, OH, OR¹, NH₂, NHCOR¹, F, CH₂OH, CH₂OR¹, a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein R¹ is hydrogen, CHO, COOR¹¹, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group or a saccharide moiety having the structure:

10

15

1

wherein Y and Z are independently NH or O; wherein k, l, r, s, t, u, v and w are each independently 0, 1 or 2; wherein R₁₀, R₁₁, R₁₂, R₁₃, R₁₄ and R₁₅ are each independently hydrogen, OH, ORⁱⁱⁱ, NH₂, NHCORⁱⁱⁱ, F, CH₂OH, CH₂ORⁱⁱⁱ, or a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein R₁₆ is hydrogen, COOH, COORⁱⁱ, CONHRⁱⁱ, a substituted or unsubstituted linear or branched chain alkyl or aryl group; wherein Rⁱⁱⁱ is hydrogen, CHO, COOR^{iv}, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group; and wherein Rⁱⁱ and R^{iv} are each independently H, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group; and

wherein each occurrence of n is independently 0-8, whereby, if for each occurrence of n, n=0, at least one occurrence of A has a different structure from other occurrences of A; and wherein the n-alkyl glycosidic moiety is either α - or β -linked to an amino acid.

20

29. The compound of claim 27 or the construct of 28, wherein each occurrence of A is independently Globo-H, fucosyl GM1, KH-1, glycophorin, Le^y, N3, Tn, 2,6-STn, (2,3)ST, or TF.

25

30. The construct of claim 28, wherein said compound has three occurrences of A comprising Tn, Globo-H and Le^y

30

31. The compound of claim 27 or the construct of claim 28, wherein at least one occurrence of A is a carbohydrate determinant having the structure:

wherein each occurrence of R' is independently hydrogen or a protecting group; and wherein each occurrence of R" is independently hydrogen or a nitrogen protecting group.

10

32. The compound of claim 27 or the construct of claim 28, wherein at least one occurrence of A is a carbohydrate determinant having the structure:

15

wherein each occurrence of R' is independently hydrogen or a protecting group; and wherein R" is hydrogen or a nitrogen protecting group.

33. A synthetic construct having the structure:

20

wherein A is a carbohydrate domain having the structure:

$$\begin{array}{c|c}
R_0 & & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_4 & \\
\hline
 & R_6 & \\
\hline
 & R_7 & \\
\hline
 & R_8 & \\
\hline
 & R_9 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_4 & \\
\hline
 & R_6 & \\
\hline
 & R_7 & \\
\hline
 & R_8 & \\
\hline
 & R_9 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_2 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_2 & \\
\hline
 & R_3 & \\
\hline
 & R_1 & \\
\hline
 & R_2 & \\
\hline
 & R_2 & \\
\hline
 & R_3 &$$

10

5

wherein a, b, c, d, e, f, g, h, i, x, y and z are independently 0, 1, 2 or 3, with the proviso that x, y and z are not simultaneously 0; wherein R₀ is hydrogen, a linear or branched chain alkyl, acyl, arylalkyl or aryl group; wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈ and R₉ are each independently hydrogen, OH, ORⁱ, NH₂, NHCORⁱ, F, CH₂OH, CH₂ORⁱ, a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein Rⁱ is hydrogen, CHO, COORⁱⁱ, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group or a saccharide moiety having the structure:

10

15

wherein Y and Z are independently NH or O; wherein k, l, r, s, t, u, v and w are each independently 0, 1 or 2; wherein R₁₀, R₁₁, R₁₂, R₁₃, R₁₄ and R₁₅ are each independently hydrogen, OH, ORⁱⁱⁱ, NH₂, NHCORⁱⁱⁱ, F, CH₂OH, CH₂ORⁱⁱⁱ, or a substituted or unsubstituted linear or branched chain alkyl, (mono-, di- or tri)hydroxyalkyl, (mono-, di- or tri)acyloxyalkyl, arylalkyl or aryl group; wherein R₁₆ is hydrogen, COOH, COORⁱⁱ, CONHRⁱⁱ, a substituted or unsubstituted linear or branched chain alkyl or aryl group; wherein Rⁱⁱⁱ is hydrogen, CHO, COOR^{iv}, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group; and wherein Rⁱⁱ and R^{iv} are each independently H, or a substituted or unsubstituted linear or branched chain alkyl, arylalkyl or aryl group;

wherein n is 0-8; wherein the carrier a lipid or protein linked directly or through a crosslinker; and wherein m is in the range of 20 to 600.

- 20 34. The construct of claim 33, wherein m is in the range of 200 to 600.
 - 35. The construct of claim 33, wherein n is 4.
- 36. The construct of claim 33, wherein the carbohydrate determinant is selected from the group consisting of Globo-H, fucosyl GM1, KH-1, glycophorin, STN, (2,3)ST, 2,6-STn, N3, Tn, TF and Le^y.
 - 37. The construct of claim 33 or 35, wherein A is a carbohydrate determinant, having the structure:

30

- wherein each occurrence of R' is independently hydrogen or a protecting group; and wherein each occurrence of R" is independently hydrogen or a nitrogen protecting group.
 - 38. The construct of claim 33 or 35, wherein A is a carbohydrate determinant, having the structure:

15

wherein each occurrence of R' is independently hydrogen or a protecting group; and wherein R" is hydrogen or a nitrogen protecting group.

20

1.

5 39. A pharmaceutical composition comprising:
a compound or construct of claim 1 or 27, and
a pharmaceutically suitable carrier.

(;

- 40. A method of treating cancer in a subject suffering therefrom comprising:

 administering to a subject a therapeutically effective amount of a compound or construct of claim 1 or 27,

 and a pharmaceutically suitable carrier.
- 41. The method of claim 40, wherein said method comprises preventing the recurrence of cancer in a subject.
 - 42. The method of claim 40 or 41, further comprising co-administering one or more immunological adjuvants.
- 20 43. The method of claim 42, wherein at least one of said one or more immunological adjuvants is a saponin adjuvant.
 - 44. The method of claim 43, wherein said saponin adjuvant is GPI-0100.
- 25 45. The method of claim 42, wherein at least one of said one or more immunological adjuvants is bacteria or liposomes.
 - 46. The method of claim 45, wherein the immunological adjuvant is Salmonella minnesota cells, bacille Calmette-Guerin or QS21.
 - 47. The method of claim 40 or 41, wherein the cancer is a solid tumor.
 - 48. The method of claim 40 or 41, wherein the subject is in clinical remission, or where the subject has been treated by surgery, has limited unresected disease.

30

- 49. A method of inducing antibodies in a subject, wherein the antibodies are capable of specifically binding with tumor cells, which comprises administering to the subject an amount of a compound or construct of claim 1 or 27 effective to induce the antibodies.
- 50. The method of claim 49, further comprising co-administering one or more immunological adjuvants.
 - 51. The method of claim 50, wherein at least one of said one or more immunological adjuvants is a saponin adjuvant.
- 15 52. The method of claim 51, wherein said saponin adjuvant is GPI-0100.
 - 53. The method of claim 49, wherein at least one of said one or more immunological adjuvants is bacteria or liposomes.
- 20 54. The method of claim 53, wherein the immunological adjuvant is Salmonella minnesota cells, bacille Calmette-Guerin or QS21.
 - 55. The method of claim 49, wherein the subject is in clinical remission, or where the subject has been treated by surgery, has limited unresected disease.

25

1/17.

Figure 1

Figure 2

1

į




Figure 3

Figure 4

Figure 5

Figure 6

m = 20-600

Figure 7

į

8/17

n = 20-600

hexasaccharide

Figure 8

Figure 9

Figure 10

Figure 11

WO 01/14395

12/17

OH
$$\begin{array}{c}
\text{NaCNBH}_{3} \\
\text{HO}
\end{array}$$

$$\begin{array}{c}
\text{NaCNBH}_{3} \\
\text{Ho}
\end{array}$$

$$\begin{array}{c}
\text{NH}
\end{array}$$

$$\begin{array}{c}
\text{NH}$$

$$\begin{array}{c}
\text{NH}
\end{array}$$

$$\begin{array}{c}
\text{NH}$$

$$\begin{array}{c}
\text{NH}
\end{array}$$

$$\begin{array}{c}
\text{NH}
\end{array}$$

$$\begin{array}{c}
\text{NH}$$

$$\begin{array}{c}
\text{NH}
\end{array}$$

Figure 12

Globo-H

R = $Fuc\alpha 1-2Gal\beta 1-3GalNAc\beta 1-3Gal\alpha 1-4Gal\beta 1-$

Fucosyl GM1

R = $Fuc\alpha 1-2Gal\beta 1-3GalNAc\beta 1-4Gal(\alpha 2-3NeuNAc)Gal\beta 1-$

Figure 13

Figure 14




Figure 15

Figure 16

Figure 17

(19) World Intellectual Property Organization International Bureau



(43) International Publication Date 1 March 2001 (01.03.2001)

PCT

(10) International Publication Number WO 01/14395 A3

(51) International Patent Classification7: C07H 15/04, A61K 31/70, A61P 35/00

(21) International Application Number: PCT/US00/22894

(22) International Filing Date: 18 August 2000 (18.08.2000)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data: 60/150,088

20 August 1999 (20.08.1999) US

- (71) Applicant: SLOAN-KETTERING INSTITUTE FOR CANCER RESEARCH [US/US]; 1275 York Avenue, New York, NY 10021 (US).
- (72) Inventors: DANISHEFSKY, Samuel, J.; 22 Brayton Street, Englewood, NJ 07631 (US). ALLEN, Jennifer, R.; 5886 North Delaware Street, Indianapolis, IN 46220 (US). RAGUPATHI, Govindaswami; 504 East 81st Street, New York, NY 10028 (US). LIVINGSTON, Philip, O.; 156 East 79th Street, New York, NY 10021 (US). WILLIAMS, Lawrence; 401 East 89th Street, Apartment 1L, New York, NY 10021 (US).

- (74) Agent: SHAIR, Karoline, K., M.; Choate, Hall & Stewart, Exchange Place, 53 State Street, Boston, MA 02109 (US).
- (81) Designated States (national): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, Cl, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- (88) Date of publication of the international search report: 7 September 2001

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

A3

(54) Title: NOVEL GLYCOCONJUGATES, GLYCOAMINO, ACIDS, INTERMEDIATES THERETO, AND USES THEREOF

(57) Abstract: The present invention provides novel n-alkenyl glycosides and glycoconjugates, n-alkyl glycoamino acids, and methods for the synthesis thereof. In another aspect, the present invention provides novel clustered glycopeptides and methods for the synthesis thereof. In still another aspect, the present invention provides methods for the treatment of cancer, preferably for the prevention of recurrence of cancer, and methods for inducing antibodies in a subject, comprising administering to a subject in need, an effective amount of any of the inventive glycoconjugates as disclosed herein.

Int tional Application No PCT/US 00/22894

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C07H15/04 A61 A61K31/70 A61P35/00 According to International Patent Classification (IPC) or to both national classification and IPC **B. FIELDS SEARCHED** Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07H A61K A61P Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practical, search terms used) CHEM ABS Data, EPO-Internal, WPI Data C. DOCUMENTS CONSIDERED TO BE RELEVANT Category ° Citation of document, with indication, where appropriate, of the relevant passages Relevant to claim No. 1,2,11, 13,39-55 Х WO 96 34005 A (SLOAN KETTERING INST CANCER) 31 October 1996 (1996-10-31) claims S. F. SLOVIN ET AL.: "Carbohydrate Χ 1,2,11, vaccines in cancer: Immunogenicity of a 13,33, fully synthetic globo H hexasaccharide 34,36, conjugate in man* 38-55 PROC. NATL. ACAD. SCI. USA, vol. 96, 1999, pages 5710-5715, XP002155262 the whole document Further documents are listed in the continuation of box C. Patent family members are listed in annex. Special categories of cited documents: T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the "A" document defining the general state of the art which is not considered to be of particular relevance nomevni earlier document but published on or after the international filing date "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such docucitation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or ments, such combination being obvious to a person skilled in the art. document published prior to the international filing date but later than the priority date claimed "&" document member of the same patent family Date of the actual completion of the international search Date of mailing of the international search report 10.04.2001 14 December 2000 Name and mailing address of the ISA Authorized officer European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016 de Nooy, A

Intr tional Application No PC (/US 00/22894

AGUPATHI, GOVINDASWAMI ET AL:	1,2,11, 13,33,
Immunization of mice with a fully	
synthetic globo H antigen results in antibodies against human cancer cells: a combined chemical-immunological approach to the fashioning of an anticancer accine" NGEW. CHEM., INT. ED. ENGL. (1997), 6(1/2), 125-128, XP002155263 Litted in the application he whole document	34,36, 38-55
AGUPATHI, GOVINDASWAMI ET AL: "A fully ynthetic globo H carbohydrate vaccine nduces a focused humoral response in rostate cancer patients: a proof of rinciple" NGEW. CHEM., INT. ED. (1999), 38(4), 63-566, XP002155264 he whole document	1,2,11, 13,33, 34,36, 38-55
ARK, TAE KYO ET AL: "Total Synthesis and roof of Structure of a Human Breast Tumor Globo-H) Antigen" . AM. CHEM. SOC. (1996), 118(46), 1488-11500, XP002155265 ited in the application age 11494, structure 64	1,2,11,
ASSALETTA, JOSE ET AL: "Glycosyl midates. Part 75. Synthesis of the exasaccharide moiety of globo H (human reast cancer) antigen" IEBIGS ANN. (1996), (9), 1417-1423, XP002155266 age 1417, structure 1b	1
. FRASER-REID ET AL.: "n-Pentenyl lycosides in organic chemistry: A contemporary example of serendipity" YNLETT, 1992, pages 927-942, XP002155267 ited in the application age 932, scheme 6 age 933, reaction compounds 37,51,52 age 937, reaction compounds 83-86 age 938, scheme 11	4,14
	o the fashioning of an anticancer accine" NGEW. CHEM., INT. ED. ENGL. (1997), 6(1/2), 125-128, XP002155263 ited in the application he whole document AGUPATHI, GOVINDASWAMI ET AL: "A fully ynthetic globo H carbohydrate vaccine nduces a focused humoral response in rostate cancer patients: a proof of rinciple" NGEW. CHEM., INT. ED. (1999), 38(4), 63-566, XP002155264 he whole document ARK, TAE KYO ET AL: "Total Synthesis and roof of Structure of a Human Breast Tumor Globo-H) Antigen" . AM. CHEM. SOC. (1996), 118(46), 1488-11500, XP002155265 ited in the application age 11494, structure 64 ASSALETTA, JOSE ET AL: "Glycosyl midates. Part 75. Synthesis of the exasaccharide moiety of globo H (human reast cancer) antigen" IEBIGS ANN. (1996), (9), 1417-1423, XP002155266 age 1417, structure 1b . FRASER-REID ET AL.: "n-Pentenyl lycosides in organic chemistry: A ontemporary example of serendipity" YNLETT, 1992, pages 927-942, XP002155267 ited in the application age 932, scheme 6 age 933, reaction compounds 37,51,52 age 937, reaction compounds 83-86 age 938, scheme 11

intr tional Application No PCT/US 00/22894

.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT alegory Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.		
U. E. UDODONG ET AL.: "A ready, convergent synthesis of the heptasaccharide GPI membrane anchor of rat brain Thy-1 glycoprotein" J. AM. CHEM. SOC., vol. 115, 1993, pages 7886-7887, XP002155268 cited in the application page 7887, scheme III	4,14		
D. CABARET, M. WAKSELMAN: "Amphiphilic liposaccharides. Synthesis and reductive cleavage of C-allyl, O-allyl, and O-butenyl glycosyl derivatives" CARBOHYDRATE RESEARCH, vol. 189, 1989, pages 341-348, XP002155269 page 341, compounds 11 and 12	3		
C. BAYLE ET AL.: "O-(3-Butenyl), a stable blocking group removable by ozonolysis" CARBOHYDRATE RESEARCH, vol. 232, 1992, pages 375-380, XP002155270 page 376, compounds 1 and 2	3		
WO 98 46246 A (HINTERMANN SAMUEL; LIVINGSTON PHILIP O (US); SLOAN KETTERING INST) 22 October 1998 (1998-10-22) figures 18-21	27		
BRODDEFALK J ET AL: "Preparation of a Glycopeptide Analogue of Type II Collagen Use of Acid Labile Protective Groups for Carbohydrate Moieties in Solid Phase Synthesis of O-Linked Glycopeptides" TETRAHEDRON LETTERS, NL, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, vol. 37, no. 17, 22 April 1996 (1996-04-22), pages 3011-3014, XP004029705 ISSN: 0040-4039 page 3011, structure 1	27		
ALLEN, JENNIFER R. ET AL: "A second generation synthesis of the MBr1 (Globo-H) breast tumor antigen: new application of the n-pentenyl glycoside method for achieving complex carbohydrate protein linkages" CHEMEUR. J. (2000), 6(8), 1366-1375,	1,2,4, 11,13, 14,16, 33-36, 38-55		

rnational application No. PCT/US 00/22894

INTERNATIONAL SEARCH REPORT

Box I Observati ns where certain claims were f und unsearchable (Continuati n f item 1 of first sheet)
This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 40-55 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
Claims Nos.: because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically: see FURTHER INFORMATION sheet PCT/ISA/210
SEE TORTHER INTORUNTION SHEEL TOTTISAY 210
3. Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).
Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)
This International Searching Authority found multiple inventions in this international application, as follows:
see additional sheet
1. As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. X No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.: See PCT/ISA/ extra sheet
Remark on Protest The additional search fees were accompanied by the applicant's protest. No protest accompanied the payment of additional search fees.

Continuation of Box I.2

Claims Nos.: 1-14 (in part), 17 (in part), 21-28 (in part), 33-35 (in part), 39-55 (in part)

Present claims 1-14, 17, 21-28, 33-35, 39-55 relate to an extremely large number of possible compounds/products/methods. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds/products/methods claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds/products/methods in which A of the first structure of claim 1 is chosen from the possibilities of claim 11, that is, A is restricted to be: Globo-H, fucosyl GM1, KH-1, glycophorin, N3, Tn, TF, STN, (2,3)ST, 2,6-STn, and Ley; and R of the first structure of claim 1 is chosen from: allyl, -CH2CH(CO2R')(NHR''), wherein R' or R'' are each independently hydrogen, protecting group, substituted or unsubstituted alkyl, a linker, aryl, peptide, protein or lipid; or NHR''', wherein R''' is a protein, peptide, or lipid linked to N directly or through a crosslinker.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-11 (in part), 13 (in full), 14 (in part), 16 (in full), 17-18 (in part), 20 (in full), 21-36 (in part), 38 (in full), 39-55 (in part)

Compounds, methods of synthesis and uses related to the first structural formula of claim 1 where A is Globo H and R is as defined in claim 1 (as limited according to article 17(2)(a) PCT).

2. Claims: 1-11 (in part), 12 (in full), 14 (in part), 15 (in full), 17-18 (in part), 19 (in full), 21-36 (in part), 37 (in full), 39-55 (in part)

Compounds, methods of synthesis and uses related to the first structural formula of claim 1 where A is fucosyl GM1 and R is as defined in claim 1 (as limited according to article 17(2)(a) PCT).

3. Claims: 1-11 (in part), 14 (in part), 17-18 (in part), 21-36 (in part), 39-55 (in part)

Compounds, methods of synthesis and uses related to the first structural formula of claim 1 where A is KH-1 and R is as defined in claim 1 (as limited according to article 17(2)(a) PCT).

4. Claims: 1-11 (in part), 14 (in part), 17-18 (in part), 21-36 (in part), 39-55 (in part)

Compounds, methods of synthesis and uses related to the first structural formula of claim 1 where A is glycophorin and R is as defined in claim 1 (as limited according to article 17(2)(a) PCT).

5. Claims: 1-11 (in part), 14 (in part), 17-18 (in part), 21-36 (in part), 39-55 (in part)

Compounds, methods of synthesis and uses related to the first structural formula of claim 1 where A is N3 and R is as defined in claim 1 (as limited according to article 17(2)(a) PCT).

6. Claims: 1-11 (in part), 14 (in part), 17-18 (in part), 21-36 (in part), 39-55 (in part)

Compounds, methods of synthesis and uses related to the first structural formula of claim 1 where A is Tn and R is as defined in claim 1 (as limited according to article 17(2)(a) PCT).

7. Claims: 1-11 (in part), 14 (in part), 17-18 (in part), 21-36 (in part), 39-55 (in part)

Compounds, methods of synthesis and uses related to the first structural formula of claim 1 where A is TF and R is as defined in claim 1 (as limited according to article 17(2)(a) PCT).

8. Claims: 1-11 (in part), 14 (in part), 17-18 (in part), 21-36 (in part), 39-55 (in part)

Compounds, methods of synthesis and uses related to the first structural formula of claim 1 where A is STN and R is as defined in claim 1 (as limited according to article 17(2)(a) PCT).

9. Claims: 1-11 (in part), 14 (in part), 17-18 (in part), 21-36 (in part), 39-55 (in part)

Compounds, methods of synthesis and uses related to the first structural formula of claim 1 where A is (2,3)ST and R is as defined in claim 1 (as limited according to article 17(2)(a) PCT).

10. Claims: 1-11 (in part), 14 (in part), 17-18 (in part), 21-36 (in part), 39-55 (in part)

Compounds, methods of synthesis and uses related to the first structural formula of claim 1 where A is 2,6-STn and R is as defined in claim 1 (as limited according to article 17(2)(a) PCT).

11. Claims: 1-11 (in part), 14 (in part), 17-18 (in part), 21-36 (in part), 39-55 (in part)

Compounds, methods of synthesis and uses related to the

first structural formula of claim 1 where A is Ley and R is as defined in claim 1 (as limited according to article 17(2)(a) PCT).

Information on patent family members

Im :tional Application No PCT/US 00/22894

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9634005	A	31-10-1996	US AU AU CA EP	5708163 A 716699 B 5672196 A 2218884 A 0823913 A	13-01-1998 02-03-2000 18-11-1996 31-10-1996 18-02-1998
			JP US US	11504337 T 5679769 A 6090789 A	20-04-1999 21-10-1997 18-07-2000
WO 9846246	Α	22-10-1998	AU EP	6779298 A 0996455 A	11-11-1998 03-05-2000